

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number  
**WO 2004/034972 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K** (74) Agent: **LOWIN, David, A.**; P.O. Box 620535, Woodside, CA 94062-0535 (US).
- (21) International Application Number: **PCT/US2003/030788** (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date:  
30 September 2003 (30.09.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/414,756 30 September 2002 (30.09.2002) US
- (71) Applicant (*for all designated States except US*): **CYTOKINETICS, INC.** [US/US]; 280 East Grand Avenue, South San Francisco, CA 94080 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BERGNES, Gustave** [US/US]; Cytokinetics, Inc., 280 East Grand Avenue, South San Francisco, CA 94080 (US). **MORGANS, David, J., Jr.** [US/US]; 781 Vista Grande Avenue, Los Altos, CA 94024 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 2004/034972 A2**

(54) Title: **COMPOUNDS, COMPOSITIONS, AND METHODS**

(57) Abstract: Compounds, compositions and methods useful for treating cellular proliferative diseases and disorders, for example, by modulating the activity of KSP, are disclosed.

## COMPOUNDS, COMPOSITIONS, AND METHODS

### CROSS-REFERENCE TO RELATED APPLICATIONS

- [001] This application claims the benefit of co-pending provisional U.S. Application Serial No. 60/414,756, filed September 30, 2002 incorporated herein by reference.

### FIELD OF THE INVENTION

- [002] This invention relates to quinazolinone-like derivatives that are inhibitors of the mitotic kinesin KSP and are useful in the treatment of cellular proliferative diseases, for example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders and inflammation.

### BACKGROUND OF THE INVENTION

- [003] The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. Disruption of the mitotic spindle can inhibit cell division, and induce cell death. Microtubules are the primary structural element of the mitotic spindle; they are the site of action of certain existing therapeutic agents used to treat cancer, such as taxanes and vinca alkaloids. Microtubules, however, exist as elements in other types of cellular structures (including tracks for intracellular transport in nerve processes). The therapeutic targeting of microtubules can, therefore, modulate processes in addition to cellular proliferation, leading to side effects that limit the usefulness of such drugs.
- [004] Improvement in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits that would be realized if the side effects associated with the administration of these agents could be reduced. Dramatic improvements in the treatment of cancer have been associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxanes, but also the camptothecin class of topoisomerase I inhibitors.
- [005] One novel anti-proliferative mechanism entails selective inhibition of mitotic kinesins, enzymes that are essential for assembly and function of the mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. See, e.g.,

Guidebook to the Cytoskeletal and Motor Proteins, Kreis and Vale, Eds., pp. 389-394 (Oxford University Press 1999). Mitotic kinesins play essential roles during all phases of mitosis. These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force that drives the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate movement of chromosomes along spindle microtubules, as well as structural changes in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death. Mitotic kinesins are attractive targets for the discovery and development of novel anti-mitotic chemotherapeutics.

[006] Among the mitotic kinesins that have been identified is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis, KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.


[007] Human KSP (also termed HsEg5) has been described [Blangy, et al., *Cell*, 83:1159-69 (1995); Whitehead, et al., *Arthritis Rheum.*, 39:1635-42 (1996); Galgio et al., *J. Cell Biol.*, 135:339-414 (1996); Blangy, et al., *J Biol. Chem.*, 272:19418-24 (1997); Blangy, et al., *Cell Motil. Cytoskeleton*, 40:174-82 (1998); Whitehead and Rattner, *J. Cell Sci.*, 111:2551-61 (1998); Kaiser, et al., *JBC* 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426], and a fragment of the KSP gene (TRIP5) has been described [Lee, et al., *Mol. Endocrinol.*, 9:243-54 (1995); GenBank accession number L40372]. *Xenopus* KSP homologs (Eg5), as well as *Drosophila* KLP61 F/KRP1 30 have been reported.

[008] Recently, certain substituted quinazolinones have been described as inhibitors of mitotic kinesins for the treatment of cellular proliferative diseases (WO 01/30768 and WO 01/98278). It is an object of the present invention to provide novel inhibitors of mitotic kinesins such as KSP (particularly human KSP).

## SUMMARY OF THE INVENTION

[009] The present invention provides compounds, compositions and methods useful in the inhibition of mitotic kinesins, particularly KSP (more particularly human KSP). The compounds can be used to treat cellular proliferative diseases and include certain pyrrolidine- and piperidine-substituted quinazolinone derivatives.

[010] In one aspect, the invention relates to one or more compounds selected from the group represented by Formula I:



Formula I

where:

U-V is chosen from  $-N(R^6)-CR^eR^f-$ ,  $-CR^eR^f-N(R^6)-$ ,  $-N(R^6)-CR^eR^f-CR^gR^h-$ ,  $-CR^eR^f-N(R^6)-CR^gR^h-$ , and  $-CR^eR^f-CR^gR^h-N(R^6)-$ ;

$R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are independently chosen from hydrogen, alkyl, aryl, aralkyl, heteroaryl, substituted alkyl, substituted aryl, substituted aralkyl, and substituted heteroaryl;

$R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are independently chosen from hydrogen, alkyl, alkoxy, halogen, cyano and substituted alkyl;


$R^5$  is chosen from alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, substituted alkyl, substituted aryl, substituted aralkyl, substituted heteroaryl and substituted heteroaralkyl; and

$R^6$  is chosen from hydrogen, acyl, alkyl, aryl, aralkyl, heteroaryl, substituted acyl, substituted alkyl, substituted aryl, substituted aralkyl and substituted heteroaryl;

including single stereoisomers, mixtures of stereoisomers, and pharmaceutically acceptable salts, solvates, and solvates of pharmaceutically acceptable salts thereof. Compounds of Formula I and pharmaceutically acceptable salts and solvates thereof are useful as active agents in the practice of the methods of treatment and in manufacture of compositions

including the pharmaceutical formulations of the invention, and may also be useful as intermediates in the synthesis of such active agents.

[011] In another aspect, the invention relates to one or more compounds selected from the group represented by Formula II:



Formula II

where:

T is a covalent bond or optionally substituted lower alkylene;

W, X and Y are independently chosen from  $-N=$ , N,  $-C=$ , CH,  $CR^i$ , O and S;

Z is chosen from  $-N=$ , N,  $-C=$ , CH,  $CR^i$  or is absent, provided that:

no more than two of W, X, Y and Z are  $-N=$ , and

W, X or Y can be O or S only when Z is absent;

$R^i$  is chosen from alkyl, alkoxy, halogen, cyano and substituted alkyl; and

$R^1$  to  $R^6$ , U and V are as defined with regard to Formula I, provided that  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  is absent where W, X, Y or Z, respectively, is  $-N=$ , O, S or absent;

including single stereoisomers, mixtures of stereoisomers, and pharmaceutically acceptable salts, solvates, and solvates of pharmaceutically acceptable salts thereof. The compounds encompassed by Formula II and pharmaceutically acceptable salts and solvates thereof will be seen to include those of Formula I; they are likewise useful as active agents in the practice of the methods of treatment and in manufacture of compositions including the pharmaceutical formulations of the invention, and may also be useful as intermediates in the synthesis of such active agents. The dashed lines in Formula II indicate that the corresponding bond may be a single bond (e.g., where X is CH) or a double bond (e.g., where X is  $-C=$ ).

[012] In one of its particular aspects the present invention pertains to a compound represented by Formula I or II, having a substituent selected from one or more of the following for  $R^a$  to  $R^h$ ,  $R^1$  to  $R^6$ , T, U, V, or W, X, Y and Z:

- U-V is  $-N(R^6)-CR^aR^f-CR^gR^h-$ ;

- $R^a$  to  $R^h$  are chosen from hydrogen, lower alkyl (particularly methyl) and substituted lower alkyl;
- $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently chosen from hydrogen, halo (particularly chloro and fluoro), lower alkyl (particularly methyl), substituted lower alkyl, lower alkoxy (particularly methoxy), and cyano;
- $R^5$  is aralkyl or substituted aralkyl (particularly benzyl or substituted benzyl; most particularly benzyl);
- $R^8$  is acyl (particularly benzoyl), aryl (particularly phenyl), substituted aryl (particularly lower alkyl-, lower alkoxy-, and/or halo-substituted phenyl), aralkyl (particularly benzyl and phenylvinyl), heteroaralkyl, oxaaralkyl (particularly phenoxy lower alkyl), oxaheteroaralkyl, substituted acyl (particularly p-toluoxy), substituted aralkyl (particularly substituted benzyl and substituted phenylvinyl), substituted heteroaralkyl, substituted oxaaralkyl (particularly substituted phenoxy lower alkyl), or substituted oxaheteroaralkyl.
- T is lower alkylene, substituted lower alkylene or a covalent bond (particularly a covalent bond); and
- W, X, Y and Z are  $-C=$  or  $-N=$  (particularly  $-C=$ ).

Other particular aspects of the invention pertain to methods and to pharmaceutical formulations employing such a compound.

[013] In one aspect, the invention relates to methods for treating cellular proliferative diseases, for disorders that can be treated by modulating KSP kinesin activity, and for inhibiting KSP kinesin by the administration of a therapeutically effective amount of a compound of Formula I or II, or a pharmaceutically acceptable salt or solvate of such compounds. Diseases and disorders that respond to therapy with compounds of the invention include cancer, hyperplasia, restenosis, cardiac hypertrophy, immune disorders and inflammation.

[014] In another aspect, the invention relates to a pharmaceutical composition containing a therapeutically effective amount of a compound of Formula I or II or a pharmaceutically acceptable salt or solvate thereof admixed with at least one pharmaceutically acceptable excipient.

[015] Yet another aspect of the invention relates to a kit having a compound, pharmaceutically acceptable salt or solvate of Formula I or II and a package insert or other labeling including directions for treating a cellular proliferative disease by administering an effective amount of the compound, salt or solvate. In one particular such aspect, the

compound, pharmaceutically acceptable salt or solvate of Formula I or II is provided as a pharmaceutical composition.

[016] In an additional aspect, the present invention provides methods of screening for compounds that will bind to a KSP kinesin, for example compounds that will displace or compete with the binding of the compounds of the invention. The methods entail combining a labeled compound of the invention, a KSP kinesin, and at least one candidate agent and determining the binding of the candidate agent to the KSP kinesin.

[017] In a further aspect, the invention provides methods of screening for modulators of KSP kinesin activity. The methods entail combining a compound of the invention, a KSP kinesin, and at least one candidate agent and determining the effect of the candidate agent on the KSP kinesin activity.

### DETAILED DESCRIPTION OF THE INVENTION

[018] The present invention provides compounds, compositions and methods useful in the inhibition of mitotic kinesins, particularly KSP (more particularly human KSP). The compounds can be used to treat cellular proliferative diseases and include certain pyrrolidine- and piperidine-substituted quinazolinone derivatives. The invention further relates to pharmaceutical formulations comprising compounds of the invention, and methods of treatment employing such compounds or compositions.

#### Definitions

[019] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise. The following abbreviations and terms have the indicated meanings throughout:

Ac	=	acetyl
Boc	=	t-butyloxy carbonyl
c-	=	cyclo
CBZ	=	carbobenzoxyl = benzyloxycarbonyl
DCM	=	dichloromethane = methylene chloride = CH <sub>2</sub> Cl <sub>2</sub>
DMF	=	N,N-dimethylformamide
DMSO	=	dimethyl sulfoxide
Et	=	ethyl
PyBroP	=	bromo-tris-pyrrolidinophosphonium hexafluorophosphate
s-	=	secondary
t-	=	tertiary
TFA	=	trifluoroacetic acid

- [020] The substituents identified as U, V, W and Y are intended to have the meanings set forth in the Summary, this Detailed Description and the Claims; they are not intended to designate the atomic elements Uranium, Vanadium, Tungsten and Yttrium.
- [021] The term "optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted alkyl" includes "alkyl" and "substituted alkyl," as defined below. It will be understood by those skilled in the art with respect to any group containing one or more substituents that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical, synthetically non-feasible and/or inherently unstable.
- [022] "Alkyl" is intended to include linear, branched, or cyclic aliphatic hydrocarbon structures and combinations thereof, which structures may be saturated or unsaturated (particularly having up to 20 carbon atoms, more particularly up to C<sub>13</sub>). Lower alkyl refers to alkyl groups of from 1 to 5 (particularly 1 to 4) carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s- and t-butyl and the like. Cycloalkyl (or carbocyclic) is a subset of alkyl and includes cyclic aliphatic hydrocarbon groups of from 3 to 13 carbon atoms. Examples of cycloalkyl groups include c-propyl, c-butyl, c-pentyl, norbornyl, adamantyl and the like. In this application, alkyl refers to alkanyl, alkenyl and alkynyl residues; it is intended to include cyclohexylmethyl, vinyl, allyl, isoprenyl and the like. Alkylene, alkenylene and alkynylene are other subsets of alkyl, referring to the same residues as alkyl, but having two points of attachment. Examples of alkylene include ethylene (-CH<sub>2</sub>CH<sub>2</sub>-), ethenylene (-CH=CH-), propylene (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), dimethylpropylene (-CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>-) and cyclohexylpropylene (-CH<sub>2</sub>CH<sub>2</sub>CH(C<sub>6</sub>H<sub>13</sub>)-). When an alkyl residue having a specific number of carbons is named, all geometric isomers of that residue having the specified number of carbons are meant to be included; thus, for example, "butyl" is meant to include n-butyl, sec-butyl, isobutyl and t-butyl; "propyl" includes n-propyl and isopropyl.
- [023] The term "alkoxy" or "alkoxyl" refers to the group -O-alkyl, particularly including from 1 to 8 carbon atoms in a straight, branched or cyclic configuration, or combinations thereof, attached to the parent structure through an oxygen. Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy, cyclohexyloxy and the like. Lower-alkoxy refers to groups containing one to five carbons.
- [024] The term "substituted alkoxy" refers to the group -O-(substituted alkyl). One particular substituted alkoxy group is "polyalkoxy" or -O-(optionally substituted alkylene)-(optionally substituted alkoxy), and includes groups such as -OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, and glycol ethers such as polyethyleneglycol and -O(CH<sub>2</sub>CH<sub>2</sub>O)<sub>x</sub>CH<sub>3</sub>, where x is an integer of about 2-20, particularly



about 2-10, and more particularly about 2-5. Another particular substituted alkoxy group is hydroxyalkoxy or  $-OCH_2(CH_2)_yOH$ , where y is an integer of about 1-10, particularly about 1-4.

[025] "Acyl" refers to groups of from 1 to 8 carbon atoms in a straight, branched or cyclic configuration, or combinations thereof, or to a hydrogen atom attached to the parent structure through a carbonyl functionality. Such groups may be saturated or unsaturated, and aliphatic or aromatic. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples include formyl, acetyl, benzoyl, propionyl, isobutyryl, t-butoxycarbonyl, benzyloxycarbonyl, aminocarbonyl, and the like. Lower-acyl refers to an acyl group containing one to five carbons. "Substituted acyl" refers to an acyl group where one or more of the hydrogens otherwise attached to a carbon, nitrogen or sulfur atom is substituted, the point of attachment to the parent moiety remaining at the carbonyl.

[026] The term "acyloxy" refers to the group  $-O$ -acyl. "Substituted acyloxy" refers to the group  $-O$ -substituted acyl.

[027] The term "amidino" refers to the group  $-C(=NH)-NH_2$ . The term "substituted amidino" refers to the formula  $-C(=NR)-NRR$  in which each R is independently selected from the group: hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aminocarbonyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, acyl, alkoxycarbonyl, sulfanyl, sulfinyl and sulfonyl, provided that at least one R is not hydrogen.

[028] The term "amino" refers to the group  $-NH_2$ . The term "substituted amino" refers to the group  $-NHR$  or  $-NRR$  where each R is independently selected from the group: optionally substituted acyl, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, sulfinyl and sulfonyl, e.g., methylamino, dimethylamino, diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino, guanidino.

[029] "Aryl" and "heteroaryl" mean a 5- or 6-membered aromatic ring or heteroaromatic ring containing 1-4 heteroatoms selected from O, N, or S; a bicyclic 9- or 10-membered aromatic ring system or heteroaromatic ring system containing 1-4 (or more) heteroatoms selected from O, N, or S; or a tricyclic 13- or 14-membered aromatic ring system or heteroaromatic ring system containing 1-4 (or more) heteroatoms selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., benzene, naphthalene, indane, tetralin, and fluorene and the 5- to 10-membered aromatic heterocyclic rings include, e.g., imidazole, pyridine, indole, thiophene, benzopyranone, thiazole, furan, benzimidazole, quinoline, isoquinoline, quinoxaline, pyrimidine, pyrazine, tetrazole and pyrazole; particularly imidazole and

imidazoline.

[030] "Aralkyl" refers to a residue in which an aryl moiety is attached to the parent structure via an alkyl residue. Examples include benzyl, phenethyl, phenylvinyl, phenylallyl and the like. "Heteroaralkyl" refers to a residue in which a heteroaryl moiety is attached to the parent structure via an alkyl residue. Examples include furanylmethyl, pyridinylmethyl, pyrimidinylethyl and the like.

[031] The term "aryloxy" refers to the group -O-aryl. Similarly, "aralkoxy" and "heteroaralkoxy" refer, respectively, to an aryl or heteroaryl moiety attached to the parent structure via an alkoxy residue.

[032] "Halogen" or "halo" refers to fluorine, chlorine, bromine or iodine (particularly fluorine, chlorine and bromine). Dihaloaryl, dihaloalkyl, trihaloaryl etc. refer to aryl and alkyl substituted with a plurality of halogens, but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl.

[033] "Heterocycle" or "heterocyclyl" means a cycloalkyl or aryl residue in which one to four of the carbons is replaced by a heteroatom such as oxygen, nitrogen or sulfur (i.e., encompassing heterocycloalkyl and heteroaryl). Examples of heterocyclyl residues that fall within the scope of the invention include imidazolyl, imidazoliny, pyrrolidinyl, pyrazolyl, pyrrolyl, indolyl, quinoliny, isoquinoliny, tetrahydroisoquinoliny, benzofuranyl, benzodioxanyl, benzodioxolyl (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazolyl, morpholiny, thiazolyl, pyridiny, pyridaziny, pyrimidinyl, thiophenyl, furanyl, oxazolyl, oxazoliny, isoxazolyl, dioxanyl, tetrahydrofuranyl and the like. "N-heterocyclyl" refers to a nitrogen-containing heterocycle as a substituent residue. Examples of N-heterocyclyl residues include 4-morpholiny, 4-thiomorpholiny, 1-piperidinyl, 1-pyrrolidinyl, 3-thiazolidinyl, piperazinyl and 4-(3,4-dihydrobenzoxazinyl). Examples of substituted heterocyclyl include 4-methyl-1-piperazinyl and 4-benzyl-1-piperidinyl.

[034] The terms "heteroaryloxy" and "heterocycloxy" refer, respectively to the groups -O-heteroaryl and -O-heterocyclyl.

[035] The term "solvate" refers to a compound (e.g., a compound of Formula I or II or a pharmaceutically acceptable salt thereof) in physical association with one or more molecules of a pharmaceutically acceptable solvent. It will be understood that phrases such as "a compound of Formula I or II or a pharmaceutically acceptable salt or solvate thereof" are intended to encompass the compound of Formula I or II, a pharmaceutically acceptable salt of the compound, a solvate of the compound, and a solvate of a pharmaceutically acceptable salt of the compound.

[036] The term "substituted" as used with regard to alkyl, aryl, aralkyl, heteroaryl and

heterocyclyl refers to an alkyl, aryl, aralkyl, heteroaryl or heterocyclyl moiety wherein one or more (up to about 5, particularly up to about 3) hydrogen atoms are replaced by a substituent independently selected from the group: optionally substituted acyl (e.g., aminocarbonyl and alkoxy carbonyl or "esters"), optionally substituted acyloxy (e.g., acid esters, carbamic acid esters, carbonic acid esters, and thiocarbonic acid esters), optionally substituted alkyl (e.g., fluoroalkyl), optionally substituted alkoxy (e.g., methoxy and methoxymethoxy), alkylenedioxy (e.g. methylenedioxy), optionally substituted amino (e.g., alkylamino, dialkylamino, carbonylamino, benzyloxycarbonylamino or "CBZ-amino", and carboxamido), optionally substituted amidino, optionally substituted aryl (e.g., phenyl and 4-methyl-phenyl or "tolyl"), optionally substituted aralkyl (e.g., benzyl), optionally substituted aryloxy (e.g., phenoxy), optionally substituted aralkoxy (e.g., benzyloxy), optionally substituted heteroaryl, optionally substituted heteroaralkyl, optionally substituted heteroaryloxy, optionally substituted heteroaralkoxy, carboxy (-COOH), cyano, halogen, hydroxy, nitro, sulfanyl, sulfinyl, sulfonyl and thio. In the compounds of Formula II where T is substituted alkylene, the term "substituted" also refers to alkylene groups where one or more (up to about 3, particularly 1) carbon atoms are replaced by a heteroatom independently selected from O, N or S, such as -CH<sub>2</sub>-S-CH<sub>2</sub>-.

[037] The term "sulfanyl" refers to the groups: -S-(optionally substituted alkyl), -S-(optionally substituted aryl), -S-(optionally substituted heteroaryl), and -S-(optionally substituted heterocyclyl).

[038] The term "sulfinyl" refers to the groups: -S(O)-H, -S(O)-(optionally substituted alkyl), -S(O)-(optionally substituted amino), -S(O)-(optionally substituted aryl), -S(O)-(optionally substituted heteroaryl), and -S(O)-(optionally substituted heterocyclyl).

[039] The term "sulfonyl" refers to the groups: -S(O<sub>2</sub>)-H, -S(O<sub>2</sub>)-(optionally substituted alkyl), -S(O<sub>2</sub>)-(optionally substituted amino), -S(O<sub>2</sub>)-(optionally substituted aryl), -S(O<sub>2</sub>)-(optionally substituted heteroaryl), -S(O<sub>2</sub>)-(optionally substituted heterocyclyl), -S(O<sub>2</sub>)-(optionally substituted alkoxy), -S(O<sub>2</sub>)-(optionally substituted aryloxy), -S(O<sub>2</sub>)-(optionally substituted heteroaryloxy), and -S(O<sub>2</sub>)-(optionally substituted heterocycliloxy).

[040] "Isomers" are different compounds that have the same molecular formula. "Stereoisomers" are isomers that differ only in the way the atoms are arranged in space. "Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The term "(.±.)" is used to designate a racemic mixture where appropriate. "Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other. The absolute stereochemistry is specified according to the Cahn-Ingold-Prelog R-S system. When

a compound is a pure enantiomer the stereochemistry at each chiral carbon may be specified by either R or S. Resolved compounds whose absolute configuration is unknown are designated (+) or (-) depending on the direction (dextro- or levorotatory) which they rotate plane polarized light at the wavelength of the sodium D line. The term "substantially pure" means having at least about 95% chemical purity with no single impurity greater than about 1%. The term "substantially optically pure" or "substantially enantiomerically pure" means having at least about 95% enantiomeric excess. The invention contemplates the use of pure enantiomers and mixtures of enantiomers, including racemic mixtures, although the use of a substantially optically pure enantiomer will generally be most suitable.

[041] "Mitotic spindle formation" refers to the organization of microtubules into bipolar structures by mitotic kinesins. "Mitotic spindle dysfunction" refers to mitotic arrest, monopolar spindle formation or mitotic spindle malformation, in which context "malformation" encompasses the splaying of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. The term "inhibit" as used with reference to mitotic spindle formation, means altering mitotic spindle formation, including decreasing spindle formation, and increasing or decreasing spindle pole separation. "Anti-mitotic" means inhibiting or having the potential to inhibit mitosis, for example, as described above.

[042] The term "pharmaceutically acceptable salts" is meant to include both acid and base addition salts. A "pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, or organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly suitable are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

[043] The term "therapeutically effective amount" or "effective amount" refers to that amount of a compound of Formula I or II that is sufficient to effect treatment, as defined below, when administered to a patient in need of such treatment. The effective amount will vary depending

upon the patient and disease condition being treated, the weight and age of the patient, the severity of the disease condition, the particular compound, pharmaceutically acceptable salt or solvate of Formula I or II chosen, the dosing regimen to be followed, timing of administration, the manner of administration and the like, all of which can readily be determined by one of ordinary skill in the art. In a particular aspect of the invention, the effective amount will be an amount sufficient to inhibit KSP kinesin activity in cells involved with the disease being treated.

[044] The term "treatment" or "treating" means any treatment of a disease in a patient, including:


- a) preventing the disease, that is, causing the clinical symptoms of the disease not to develop;
- b) inhibiting the disease, that is, slowing or arresting the development of clinical symptoms; and/or
- c) relieving the disease, that is, causing the regression of clinical symptoms.

[045] A "patient" for the purposes of the present invention includes humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In a particular embodiment the patient is a mammal, most particularly the patient is human.

#### COMPOUNDS OF THE PRESENT INVENTION

[046] The present invention provides certain quinazolinone derivatives. The compounds are inhibitors of one or more mitotic kinesins. The present invention capitalizes on the finding that perturbation of mitotic kinesin function causes malformation or dysfunction of mitotic spindles, frequently resulting in cell cycle arrest and cell death.

[047] Accordingly, the present invention relates to one or more compounds selected from the group represented by Formula I:



U-V is chosen from  $-N(R^6)-CR^aR^f$ -,  $-CR^aR^f-N(R^6)-$ ,  $-N(R^6)-CR^aR^f-CR^aR^h$ -,  
 $-CR^aR^f-N(R^6)-CR^aR^h$ -, and  $-CR^aR^f-CR^aR^h-N(R^6)-$ ;

$R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are independently chosen from hydrogen, alkyl, aryl, aralkyl, heteroaryl, substituted alkyl, substituted aryl, substituted aralkyl and substituted heteroaryl;


$R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are independently chosen from hydrogen, alkyl, alkoxy, halogen, cyano and substituted alkyl;

$R^5$  is chosen from alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, substituted alkyl, substituted aryl, substituted aralkyl, substituted heteroaryl and substituted heteroaralkyl; and

$R^6$  is chosen from hydrogen, acyl, alkyl, aryl, aralkyl, heteroaryl, substituted acyl, substituted alkyl, substituted aryl, substituted aralkyl and substituted heteroaryl;

including single stereoisomers, mixtures of stereoisomers, and pharmaceutically acceptable salts, solvates, and solvates of pharmaceutically acceptable salts thereof.

[048] In another aspect, the invention relates to one or more compounds selected from the group represented by Formula II:



Formula II

where:

T is a covalent bond or optionally substituted lower alkylene;

W, X and Y are independently chosen from  $-N=$ , N,  $-C=$ , CH,  $CR^i$ , O and S;

Z is chosen from  $-N=$ , N,  $-C=$ , CH,  $CR^i$  or is absent, provided that:

no more than two of W, X, Y and Z are  $-N=$ , and

W, X or Y can be O or S only when Z is absent;

$R^i$  is chosen from alkyl, alkoxy, halogen, cyano and substituted alkyl; and

$R^1$  to  $R^6$ , U and V are as defined with regard to Formula I, provided that  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  is absent where W, X, Y or Z, respectively, is  $-N=$ , O, S or absent;

including single stereoisomers, mixtures of stereoisomers, and the pharmaceutically acceptable salts, solvates, and solvates of pharmaceutically acceptable salts thereof. The


compounds encompassed by Formula II will be seen to include those of Formula I; they are likewise useful as active agents in the practice of the methods of treatment and in manufacture of compositions including the pharmaceutical formulations of the invention, and may also be useful as intermediates in the synthesis of such active agents. The dashed lines in Formula II indicate that the corresponding bond may be a single bond (e.g., where X is CH) or a double bond (e.g., where X is  $-C=$ ). For the sake of simplicity in the following description and claims, substituents T, U, W, X, Y and Z will not be discussed in connection with certain compounds falling within the scope of Formula I.

[049] Many of the compounds described herein contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that such compounds include both E and Z geometric isomers. All tautomeric forms are also intended to be included. The present invention is meant to include all such possible isomers, including racemic mixtures, intermediate mixtures, optically pure forms, substantially optically pure forms, enantiomerically pure forms, and substantially enantiomerically pure forms.

#### Nomenclature

[050] The compounds of Formula I and II can be named and numbered (e.g., using AutoNom version 2.1 in ISIS-DRAW or ChemDraw) as described below.


[051] For example, the compound of Formula IA:



Formula IA

i.e., the compound according to Formula I where U-V is  $-N(R^6)-CR^eR^f-$ ;  $R^1$ ,  $R^2$ ,  $R^4$ , and  $R^a$  to  $R^f$  are H;  $R^3$  is chloro;  $R^5$  is benzyl; and  $R^6$  is p-methyl-benzoyl, can be named 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3H-quinazolin-4-one.


[052] The compound of Formula IB:



Formula IB

i.e., the compound according to Formula I where U-V is  $-N(R^6)-CR^eR^f-CR^gR^h-$ ;  $R^1$ ,  $R^2$ ,  $R^4$  and  $R^a$  to  $R^h$  are H;  $R^3$  is chloro;  $R^5$  is benzyl; and  $R^6$  is p-methyl-benzoyl, can be named 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3H-quinazolin-4-one.

[053] The compound of Formula IC:




Formula IC

i.e., the compound according to Formula I where U-V is  $-CR^eR^f-N(R^6)-CR^gR^h-$ ;  $R^1$ ,  $R^4$ , and  $R^a$  to  $R^h$  are H;  $R^2$  and  $R^3$  are methoxy;  $R^5$  is benzyl; and  $R^6$  is p-methyl-benzyl, can be named 3-benzyl-6,7-dimethoxy-2-[1-(4-methyl-benzyl)-piperidin-3-yl]-3H-quinazolin-4-one.




[054] The compound of Formula ID:



Formula ID

i.e., the compound according to Formula I where U-V is  $-\text{CR}^{\text{a}}\text{R}^{\text{f}}-\text{CR}^{\text{g}}\text{R}^{\text{h}}-\text{N}(\text{R}^{\text{b}})-$ ;  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^{\text{b}}$  and  $\text{R}^{\text{d}}$  to  $\text{R}^{\text{h}}$  are H;  $\text{R}^3$  is chloro;  $\text{R}^5$  is methyl;  $\text{R}^{\text{a}}$  is phenethyl;  $\text{R}^{\text{c}}$  is diethylamino-ethyl; and  $\text{R}^{\text{e}}$  is p-methyl-benzoyl, can be named 7-chloro-2-[2-(2-diethylamino-ethyl)-1-(4-methyl-benzoyl)-3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one.

[055] The compound of Formula IIA:




Formula IIA

i.e., the compound according to Formula II where T is methylene; U-V is  $-\text{N}(\text{R}^{\text{b}})-\text{CR}^{\text{a}}\text{R}^{\text{f}}-$ ; W, Y and Z are  $-\text{C}=\text{C}-$ ; X is  $-\text{N}=\text{N}-$ ;  $\text{R}^1, \text{R}^4$  and  $\text{R}^{\text{a}}$  to  $\text{R}^{\text{f}}$  are H;  $\text{R}^2$  is absent;  $\text{R}^3$  is chloro;  $\text{R}^5$  is benzyl; and  $\text{R}^{\text{e}}$  is p-methyl-benzoyl, can be named 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-ylmethyl]-3H-pyrido[4,3-d]pyrimidin-4-one.

[056]

The compound of Formula IIB:




Formula IIB

i.e., the compound according to Formula I where the dashed line between W and X is a single bond, T is 3-propionyl; U-V is  $-N(R^6)-CR^eR^f-CR^gR^h-$ ; W is N; X is  $CR^i$  where  $R^i$  is methyl; Y is CH; Z is =N-;  $R^1$ ,  $R^4$  and  $R^a$  to  $R^h$  are H;  $R^2$  is methyl;  $R^3$  is absent;  $R^5$  is benzyl; and  $R^6$  is p-methyl-benzoyl, can be named 3-benzyl-6,6-dimethyl-2-{3-[1-(4-methyl-benzoyl)-piperidin-2-yl]-propionyl}-5,6-dihydro-3H-pteridin-4-one.

[057]


The compound of Formula IIC:



Formula IIC

i.e., the compound according to Formula II where the dashed lines are single bonds, T is absent; U-V is  $-CR^eR^f-N(R^6)-CR^gR^h-$ ; Z,  $R^2$  and  $R^4$  are absent; W and Y are CH; X is S;  $R^1$ ,  $R^3$ , and  $R^a$ ,  $R^b$ , and  $R^d$  to  $R^h$  are H;  $R^c$  is isopropyl;  $R^5$  is benzyl; and  $R^6$  is p-methyl-benzyl, can be named 3-benzyl-2-[5-isopropyl-1-(4-methyl-benzyl)-piperidin-3-yl]-5,7-dihydro-3H-thieno[3,4-d]pyrimidin-4-one.

[058] The compound of Formula IID:



Formula IID

i.e., the compound according to Formula II where T is aminomethylene; U-V is  $-\text{CR}^a\text{R}^f-\text{CR}^g\text{R}^h-\text{N}(\text{R}^6)-$ ;  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^b$  and  $\text{R}^d$  to  $\text{R}^h$  are H;  $\text{R}^3$  is chloro;  $\text{R}^5$  is methyl;  $\text{R}^a$  is benzyl;  $\text{R}^c$  is diethylamino-ethyl; and  $\text{R}^6$  is p-carbonyl-benzamide, can be named 4-[3-benzyl-4-[(7-chloro-3-methyl-4-oxo-3,4-dihydro-quinazolin-2-yl)methyl]-amino]-2-(2-diethylamino-ethyl)-piperidine-1-carbonyl-benzamide.

#### Synthetic Reaction Parameters

[059] The terms "solvent", "inert organic solvent" or "inert solvent" mean a solvent inert under the conditions of the reaction being described in conjunction therewith [including, for example, benzene, toluene, acetonitrile, tetrahydrofuran ("THF"), dimethylformamide ("DMF"), chloroform, methylene chloride (or dichloromethane), diethyl ether, methanol, pyridine and the like]. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert organic solvents.

[060] The term "q.s." means adding a quantity sufficient to achieve a stated function, e.g., to bring a solution to the desired volume (i.e., 100%).

[061] Isolation and purification of the compounds and intermediates described herein can be effected, if desired, by any suitable separation or purification procedure such as, for example, filtration, extraction, crystallization, column chromatography, thin-layer chromatography or thick-layer chromatography, or a combination of these procedures. Specific illustrations of suitable separation and isolation procedures can be had by reference to the examples hereinbelow. However, other equivalent separation or isolation procedures can, of course, also be used.

- [062] When desired, the (R)- and (S)-isomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallisation; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallisation, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. For example, a compound of Formula I or II can be dissolved in a lower alkanol and placed on a Chiralpak AD (205 x 20 mm) column (Chiral Technologies, Inc.) conditioned for 60 min at 70% EtOAc in Hexane. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation.


#### SYNTHESIS OF THE COMPOUNDS OF FORMULA I AND II

##### Brief Description Of Reaction Schemes

- [063] Syntheses of the compounds of Formula I and II are described below with reference to Reaction Schemes 1 and 2.
- [064] Reaction Scheme 1 illustrates a synthesis of the compounds of Formulae I and II [excluding the 2-(piperidin-2-yl)-3*H*-quinazolin-4-ones]. The nitrogen atom of U-V is protected until cyclization, followed by de-protection and derivatization to provide R<sup>6</sup> substituents other than hydrogen.
- [065] Reaction Scheme 2 depicts a synthesis for the compounds of Formulae I and II [including 2-(piperidin-2-yl)-3*H*-quinazolin-4-ones], illustrating the synthesis of compounds where R<sup>6</sup> is acyl.
- [066] It will be appreciated by those skilled in the art that one or more of the reactants, steps and/or conditions described with reference to Reaction Schemes 1 and 2 may require adjustment to accommodate non-hydrogen substituents, e.g., at R<sup>1</sup> to R<sup>6</sup> and R<sup>a</sup> to R<sup>h</sup>.

Starting Materials

[067] The N-protected pyrrolidine and piperidine dicarboxylic acids of Formula 101 (e.g., pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester), the anthranilic acids of Formula 102 (e.g., 4-chloro-anthranilic acid) and other reactants are commercially available, e.g., from Aldrich Chemical Company, Milwaukee, WI or may be readily prepared by those skilled in the art using commonly employed synthetic methodology.

**Reaction Scheme 1**

- [068] Referring to Reaction Scheme 1, the  $R^6$  component of "U-V" is not specifically shown. In Formulae 102 to 106 the  $R^6$  moiety is a protecting group. In the end products of Step 5 (Formula I or II)  $R^6$  is as defined, initially being hydrogen upon de-protection of U-V and optionally being further derivatized. Products where  $R^6$  is aryl or heteroaryl can be obtained by starting with the corresponding arylated compound of Formula 101 or by a palladium catalyzed arylation (e.g., as described by Wolfe, J.P.; Tomori, H; Sadighi, J.P.; Yin, J.; and Buchwald, S.L., J. Org. Chem., 2000, 65, 1158-1174) of a de-protected product of Step 5 where  $R^6$  is hydrogen.
- [069] Preparation of Formula 103 Referring to Reaction Scheme 1, Step 1, a solution is made of an optionally substituted o-amino alicyclic, heterocyclic or (hetero)aryl acid of Formula 102 (such as anthranilic acid), a slight molar excess of an N-protected pyrrolidine- or piperidine-dicarboxylic acid of Formula 101 [where U'-V' is  $-N(R^6)-CR^eR^f$ ,  $-CR^eR^f-N(R^6)-$ ,  $-CR^eR^f-N(R^6)-CR^gR^h$  or  $-CR^eR^f-CR^gR^h-N(R^6)-$ , and  $R^6$  is an amino-protecting group, such as Fmoc and Boc] and a slight molar excess of PyBroP dissolved in an organic solvent (e.g., diisopropylethylamine in pyridine). The solution is stirred for 12 to 20 hours at room temperature to afford the corresponding compound of Formula 103, which is conventionally isolated and purified.
- [070] Preparation of Formula 104 Referring to Reaction Scheme 1, Step 2, a compound of Formula 103 is dissolved in an organic solvent (e.g., a lower alkanol such as ethanol) and treated with an aqueous hydroxide (e.g., NaOH aq. solution). The mixture is stirred at room temperature for 12 to 20 hours and then concentrated under vacuum. The residue is washed (e.g., with saturated NaCl and concentrated phosphoric acid) and then extracted (e.g., with dichloromethane). Conventional isolation and purification affords the corresponding acid of Formula 104.
- [071] Preparation of Formula 105 Referring to Reaction Scheme 1, Step 3, to a solution of Formula 104 in an organic solvent (e.g., dichloromethane) is added large molar excesses of EDC and diisopropylethylamine. After stirring at room temperature for 1 to 2 hours, a large molar excess of an amine of the formula  $H_2N-R^5$  (such as benzylamine) is added and the reaction is stirred for an additional 24 to 60 hours. The mixture is washed, dried and isolated to afford the corresponding di-carbamoyl compound of Formula 105, and may include a small amount of the corresponding cyclized compound of Formula 106, which is taken forward without further purification.
- [072] Preparation of Formula 106 Referring to Reaction Scheme 1, Step 4, a compound of Formula 105 (optionally in the presence of a compound of Formula 106) is stirred in ethylene


glycol with  $K_2CO_3$  at  $120^\circ C$  for 12 to 20 hours, allowed to cool to room temperature, and is extracted (e.g., with dichloromethane). The organic fractions are conventionally isolated and purified to afford the pure cyclized compound of Formula 106.

[073]      Preparation of Formula I or II where  $R^6$  is Hydrogen      Referring to Reaction Scheme 1, Step 5, a solution of Formula 106 where  $R^6$  of U'-V' is a Boc group, is treated with a 1:1 mixture of TFA/dichloromethane for 30 minutes to 2 hours at room temperature. The solution is concentrated under vacuum and partitioned (e.g., between dichloromethane and saturated  $NaHCO_3$ ). The aqueous layer is extracted (e.g., with dichloromethane) and the combined organic layers are conventionally isolated to afford the corresponding de-protected pyrrolidinyl- or piperidinyl product of Formula I or II, which can be purified or taken forward without further purification.

[074]      Preparation of Formula I or II where  $R^6$  is Acyl      To a solution of Formula I or II where  $R^6$  is hydrogen in an organic solvent (e.g., dichloromethane) is added large molar excesses of an acyl halide (e.g., toluoyl chloride) and diisopropylethylamine. The mixture is stirred for 30 minutes to 2 hours and then partitioned (e.g., between saturated  $NaHCO_3$  and ethyl acetate). Conventional isolation and purification of the organic phase affords the corresponding compound of Formula I or II where  $R^6$  is acyl.

[075]      Preparation of Formula I where  $R^6$  is Optionally Substituted Alkyl, Optionally Substituted Aalkyl or Optionally Substituted Heteroaralkyl      To a solution of Formula I or II where  $R^6$  is hydrogen in an organic solvent (e.g., dichloromethane) is added a slight molar excess of an aldehyde comprising  $R^{6*}$  (i.e., a compound having formula  $R^{6*}-CHO$  where  $R^{6*}-CH_2-$  is equivalent to  $R^6$  and  $R^6$  is as described above; e.g., *p*-tolualdehyde) and a large molar excess of  $NaHB(OAc)_3$ . The mixture is stirred for 30 minutes to 1 hour at room temperature. The reaction is quenched with saturated  $NaHCO_3$  and the aqueous phase extracted (e.g., with dichloromethane). Conventional isolation and purification of the organic phase affords the corresponding compound of Formula I or II where  $R^6$  is optionally substituted-alkyl, -aralkyl or -heteroaralkyl.

## Reaction Scheme 2



[076] Referring to Reaction Scheme 2, in Formula 201 the R<sup>6</sup> moiety of "U-V" is hydrogen. In Formulae 202 through 205 and in Formulae I and II, the "N-Acyl" moiety represents the -N(R<sup>6</sup>)-portion of "U-V" where R<sup>6</sup> is Acyl.



- [077]        Preparation of Formula 202 Referring to Reaction Scheme 2, Step 1, a slight molar excess of an acyl halide (e.g., *p*-toluoyl chloride) is slowly added, portionwise, to a 0°C solution of a pyrrolidine- or piperidine-carboxylic acid of Formula 201 in 2M NaOH. The mixture is stirred for an additional 30 minutes to 2 hours at room temperature, cooled to 0°C, and partitioned (e.g., between 1N HCl and dichloromethane). The organic layer is isolated and purified to afford the corresponding N-acyl-pyrrolidine- or -piperidine-carboxylic acid of Formula 202.
- [078]        Preparation of Formula 203 Referring to Reaction Scheme 2, Step 2, a compound of Formula 102 is combined with a slight molar excess of a compound of Formula 202, and one-fifth molar equivalents each of PyBroP and diisopropylethylamine in DMF. The reaction takes place at -15°C with stirring for 12 to 20 hours, allowing the temperature to rise to room temperature. The product is isolated and purified to afford the corresponding compound of Formula 203.
- [079]        Preparation of Formula 205 Referring to Reaction Scheme 2, Steps 3 and 4, a mixture of a compound of Formula 203 and 1M NaOH in a mixed solvent (e.g., methanol and dioxane) is stirred at room temperature for 5 to 10 hours. The solvents are evaporated and the residue is partitioned (e.g., between dichloromethane, saturated NaCl, and concentrated HCl). The layers are separated and the organic layer dried and concentrated under vacuum to afford the corresponding compound of Formula 204; which crude product is carried forward without further purification. The crude product is redissolved (e.g., in dichloromethane) and treated with a large molar excess of both EDC and diisopropylethylamine. After stirring for 10 to 30 minutes, a large molar excess of an amine of the formula  $H_2N-R^5$  (such as benzylamine) is added and the mixture stirred for 12 to 20 hours at room temperature. The mixture is washed, dried and isolated to afford the corresponding crude di-carbamoyl compound of Formula 205, which can be taken forward without further purification.
- [080]        Preparation of Formula I or II Referring to Reaction Scheme 2, Step 5, a mixture of a compound of Formula 205 and a large molar excess of  $K_2CO_3$  in a suitable solvent (e.g., ethylene glycol) is stirred at 120°C for 12 to 20 hours. The solution is cooled to room temperature, diluted with water, and extracted (e.g., with dichloromethane and ethyl acetate). The combined organic layers are washed, isolated and purified to afford the corresponding 1-(acyl)-pyrrolidinyl- or 1-(acyl)-piperidinyl- product of Formula I or II.
- [081]        Compounds prepared by the above-described process of the invention may be identified by the presence of a detectable amount of Formulae 105 or 205, or a protected

precursor of Formula 106. While it is well known that pharmaceuticals must meet pharmacopoeia standards before approval and/or marketing, and that synthetic reagents (such as benzylamine, ethylene glycol or NaOH) and precursors (such as Formulae 105, 106 and 205) should not exceed the limits prescribed by pharmacopoeia standards, final compounds prepared by a process of the present invention may have minor, but detectable, amounts of such materials present, for example at levels in the range of 95% purity with no single impurity greater than 1%. These levels can be detected, e.g., by emission spectroscopy. It is important to monitor the purity of pharmaceutical compounds for the presence of such materials, which presence is additionally disclosed as a method of detecting use of a process of the invention.

#### Particular Optional Processes and Last Steps

- [082] A compound of Formula 205 is cyclized under basic conditions to afford a corresponding, optionally protected 2-[1-(acyl)-pyrrolidinyl]-3*H*-quinazolin-4-one or 2-[1-(acyl)-piperidinyl]-3*H*-quinazolin-4-one of Formula I.
- [083] A protected precursor to Formula I or II (e.g., Formula 106 where R<sup>6</sup> is NHBoc) is de-protected with in TFA in a suitable solvent to afford the corresponding de-protected compound of Formula I or II.
- [084] A compound of Formula I or II where R<sup>6</sup> is hydrogen is contacted with an alkyl or aryl halide, aldehyde or acid chloride in a suitable solvent to afford the corresponding compound of Formula I or II where R<sup>6</sup> is other than hydrogen.
- [085] A racemic mixture of isomers of a compound of Formula I or II is placed on a chromatography column and separated into (R)- and (S)- enantiomers.
- [086] A compound of Formula I or II is contacted with a pharmaceutically acceptable acid to form the corresponding acid addition salt.
- [087] A pharmaceutically acceptable acid addition salt of Formula I or II is contacted with a base to form the corresponding free base of Formula I or II.

#### Particular Compounds

- [088] Particular embodiments of the invention include or employ the compounds of Formulae I and II having the following combinations and permutations of substituent groups (indented/sub-grouped, respectively, in increasing order of particularity). These are presented in support of the appended claims as well as combinations and permutations of substituent groups that may, for the sake of brevity, not be specifically claimed but should be appreciated as encompassed by the teachings of the present disclosure. In that regard, the below-described subsets for each substituent (sometimes referenced by paragraph number) are

intended to apply to that substituent alone or in combination with one, several, or all of the described subsets for the other substituents.

[089] W, X, Y and Z are independently chosen from  $-C=$  and  $-N=$ .

- W, X, Y and Z are  $-C=$ .

[090]  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently chosen from hydrogen, halo, lower alkyl, substituted lower alkyl, lower alkoxy, and cyano.

- $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, chloro, fluoro, methyl, methoxy, cyano or substituted lower alkyl.
  - $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, chloro, fluoro, methyl, methoxy or cyano.
- Where three or four of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
  - $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
    - Where halo is chloro.
      - Where  $R^3$  is hydrogen or chloro.
        - Where  $R^3$  is chloro.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.

[091]  $R^5$  is optionally substituted aralkyl.

- $R^5$  is benzyl or substituted benzyl.
  - $R^5$  is benzyl.

[092] T is optionally substituted lower alkylene or is a covalent bond (i.e., absent).

- T is a covalent bond,  $C_1$  to  $C_4$  alkylene or  $C_1$  to  $C_4$  alkylene substituted with halo or oxo.
  - T is a covalent bond or  $C_1$  to  $C_4$  alkylene.
    - T is a covalent bond.
- Where T is alkylene having a carbon substituted by a heteroatom, the heteroatom is not bound directly to the bicyclic structure.
  - T is aminoalkylene or amidoalkylene.
- T is alkylene or alkylene substituted with halo or oxo.

[093]  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are independently chosen from hydrogen, lower alkyl (especially methyl), and substituted lower alkyl.

- Where no more than one of  $R^a$  to  $R^h$  is other than hydrogen.
  - Where all of  $R^a$  to  $R^h$  are hydrogen.

[094] U-V is  $-N(R^6)-CR^eR^f-CR^gR^h-$ ,  $-CR^eR^f-N(R^6)-CR^gR^h-$ , or  $-CR^eR^f-CR^gR^h-N(R^6)-$ .

- U-V is  $-N(R^6)-CR^eR^f-CR^gR^h-$  or  $-CR^eR^f-N(R^6)-CR^gR^h-$ .

- Where  $R^6$  is optionally substituted aralkyl or optionally substituted acyl.
  - Where  $R^6$  is optionally substituted acyl.
    - Where  $R^6$  is p-methyl-benzoyl.
- U-V is  $-N(R^6)-CR^aR^f-CR^gR^h-$ .
  - Where  $R^6$  is optionally substituted aralkyl or optionally substituted acyl.
    - Where  $R^6$  is optionally substituted acyl.
      - Where  $R^6$  is p-methyl-benzoyl.

[095]

Illustrative of the suitable combinations and permutations of particular substituents are the compounds, pharmaceutically acceptable salts and solvates where T is a covalent bond, U-V is  $-N(R^6)-CR^aR^f-CR^gR^h-$ , and one or more of W, X, Y and Z,  $R^1$  to  $R^6$ , and  $R^a$  to  $R^h$  is/are as described in paragraphs 089 to 094 above, such as:

- Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
  - Where  $R^3$  is chloro.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
- Where  $R^5$  is optionally substituted benzyl.
  - $R^5$  is benzyl.
- Where all but one of  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are hydrogen, and the remaining member is chosen from hydrogen, lower alkyl (especially methyl) and substituted lower alkyl.
  - All of  $R^a$  to  $R^h$  are hydrogen.
- Where  $R^6$  is optionally substituted acyl.
  - $R^6$  is p-methyl-benzoyl.
    - Where all but one of  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are hydrogen, and the remaining member is chosen from hydrogen, lower alkyl (especially methyl) and substituted lower alkyl.
      - All of  $R^a$  to  $R^h$  are hydrogen.
        - Where  $R^5$  is optionally substituted benzyl.
          - $R^5$  is benzyl.
            - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
              - Where  $R^3$  is chloro.
              - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.

- Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
  - Where  $R^3$  is chloro.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
- Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
  - Where  $R^3$  is chloro.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
- Where  $R^5$  is optionally substituted benzyl.
  - $R^5$  is benzyl.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
      - Where  $R^3$  is chloro.
      - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
      - Where  $R^3$  is chloro.
      - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
    - Where  $R^3$  is chloro.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
  - Where  $R^5$  is optionally substituted benzyl.
    - $R^5$  is benzyl.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
      - Where  $R^3$  is chloro.
      - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.

- Where R<sup>3</sup> is chloro.
- Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.
- Where all but one of R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>d</sup>, R<sup>e</sup>, R<sup>f</sup>, R<sup>g</sup> and R<sup>h</sup> are hydrogen, and the remaining member is chosen from hydrogen, lower alkyl (especially methyl) and substituted lower alkyl.
  - All of R<sup>a</sup> to R<sup>h</sup> are hydrogen.
    - Where R<sup>5</sup> is optionally substituted benzyl.
      - R<sup>5</sup> is benzyl.
        - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
          - Where R<sup>3</sup> is chloro.
          - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.
        - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
          - Where R<sup>3</sup> is chloro.
          - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.
      - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
        - Where R<sup>3</sup> is chloro.
        - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.
    - Where R<sup>5</sup> is optionally substituted benzyl.
      - R<sup>5</sup> is benzyl.
        - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
          - Where R<sup>3</sup> is chloro.
          - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.
        - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
          - Where R<sup>3</sup> is chloro.
          - Where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen.

- Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
  - Where  $R^3$  is chloro.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
- Where  $R^5$  is optionally substituted benzyl.
  - $R^5$  is benzyl.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
      - Where  $R^3$  is chloro.
      - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
      - Where  $R^3$  is chloro.
      - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.
  - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen or three of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen and the fourth is halo, methoxy, methyl, or cyano.
    - Where  $R^3$  is chloro.
    - Where  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are hydrogen.

Thus, the compounds where T is a covalent bond and U-V is  $-N(R^6)-CR^aR^f-CR^gR^h-$ , including those where the above-described groupings and sub-groups of substituents are taken individually and/or combined together as illustrated with regard to those compounds where  $R^6$  is optionally substituted acyl, are particularly suitable for practice of the present invention.

[096] One group of compounds, pharmaceutically acceptable salts and solvates thereof, compositions including pharmaceutical formulations, and methods of manufacture and use of the present invention are those wherein the compound of Formula I or II is selected from:

- 3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-pyrrolidin-2-yl]-3H-quinazolin-4-one;
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3H-quinazolin-4-one;
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3H-quinazolin-4-one;
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-piperidin-3-yl]-3H-quinazolin-4-one;
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3H-quinazolin-4-one;
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-piperidin-4-yl]-3H-quinazolin-4-one; and
  - 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-4-yl]-3H-quinazolin-4-one,
- especially the (R)- enantiomers thereof.

[097] A particular group of compounds, pharmaceutically acceptable salts and solvates thereof, compositions including pharmaceutical formulations, and methods of manufacture and use of the present invention are those wherein the compound of Formula I or II is selected from:

- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one;
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one;
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3*H*-quinazolin-4-one; and
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-4-yl]-3*H*-quinazolin-4-one, especially the (R)- enantiomers thereof.

[098] Another particular group of compounds, pharmaceutically acceptable salts and solvates thereof, compositions including pharmaceutical formulations, and methods of manufacture and use of the present invention are those wherein the compound of Formula I or II is selected from:

- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one; and
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3*H*-quinazolin-4-one, especially the (R)- enantiomers thereof.

#### UTILITY, TESTING AND ADMINISTRATION

##### General Utility

[099] The compounds of the invention find use in a variety of applications, including as therapeutic active agents, in the practice of the methods of treatment, in compositions, particularly pharmaceutical formulations and in methods for the manufacture of pharmaceutical formulations, and as intermediates in the synthesis of such therapeutic active agents.

[0100] As will be appreciated by those in the art, mitosis can be altered in a variety of ways; that is, one can affect mitosis either by increasing, decreasing or otherwise interfering with the activity of a component in the mitotic pathway. Stated differently, mitosis can be affected (e.g., disrupted) by disturbing equilibrium, either by inhibiting or activating certain mitotic components. Similar approaches can be used to alter meiosis.

[0101] The compounds of the invention can be used to inhibit mitotic spindle formation. Such inhibition may take the form of lessening a mitotic kinesin's organization of microtubules into bipolar structures, increasing or decreasing spindle pole separation, and/or inducing mitotic spindle dysfunction. In particular, the compounds of the invention are useful to bind to and/or inhibit the activity of a mitotic kinesin, KSP, especially human KSP, although KSP kinesins from other organisms may also be used. Also included within the definition of the term "KSP"



for these purposes are variants and/or fragments of KSP. See, U.S. Patent No. 6,437,115. While other mitotic kinesins may be used in the present invention, the compounds of the invention have been shown to have specificity for KSP. Contacting a compound of the invention with a KSP kinesin, particularly human KSP kinesin, can lead to diminished KSP-mediated ATP hydrolysis activity and/or diminished KSP-mediated mitotic spindle formation activity. Meiotic spindles can be similarly disrupted.

[0102] In another embodiment, the compounds of the invention can be used to modulate one or more other human mitotic kinesins, in addition to inhibiting KSP, including: HSET (see, U.S. Patent No. 6,361,993); MCAK (see, U.S. Patent No. 6,331,424); CENP-E (see, PCT Publication No. WO 99/13061); Kif4 (see, U.S. Patent No. 6,440,684); MKLP1 (see, U.S. Patent No. 6,448,025); Kif15 (see, U.S. Patent No. 6,355,466); Kid (see, U.S. Patent No. 6,387,644); Mpp1, CMKrp, Kinl-3 (see, U.S. Patent No. 6,461,855); Kip3a (see, PCT Publication No. WO 01/96593); Kip3d (see, U.S. Patent No. 6,492,151); and RabK6.

[0103] Therapeutic uses facilitated by the mitotic kinesin-inhibitory activity of the compounds of the present invention include the treatment of disorders associated with cell proliferation. Particular disease states that can be treated by the methods, pharmaceutical formulations, and compounds provided herein include, but are not limited to, cancer (further discussed below), autoimmune disease, arthritis, graft rejection, inflammatory bowel disease, proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. In one embodiment, the invention includes application to cells or individuals afflicted or impending affliction with any one of these disorders or states.

[0104] The compounds, pharmaceutical formulations and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that can be treated include, but are not limited to:

- Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma;
- Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma;
- Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's

- sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma);
- Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma);
  - Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma;
  - Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors;
  - Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma);
  - Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma);
  - Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma];
  - Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and
  - Adrenal glands: neuroblastoma.

As used herein, treatment of cancer includes treatment of cancerous cells, including cells afflicted by any one of the above-identified conditions.

- [0105] Another useful aspect of the invention is a kit having a compound, salt or solvate of Formula I or II and a package insert or other labeling including directions treating a cellular proliferative disease by administering an effective amount of the compound, salt or solvate. The compound, salt or solvate of Formula I or II in the kits of the invention is particularly provided as one or more doses for a course of treatment for a cellular proliferative disease, each dose being a pharmaceutical formulation including a pharmaceutically accepted excipient and a compound, salt or solvate of Formula I or II.

#### Testing

- [0106] To assay activity, generally, either KSP or a compound according to the invention is non-diffusably bound to an insoluble support having isolated sample receiving areas. The insoluble support can be made of any material to which the compounds can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports can be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the compound is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the compound and is nondiffusable. Particular methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas can then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

- [0107] The compounds of the invention can be used on their own to modulate the activity of a mitotic kinesin, particularly KSP. In this embodiment, a compound of the invention is combined with KSP and the activity of KSP is assayed. Measurable kinesin activities include the ability to affect ATP hydrolysis; microtubule binding; gliding and polymerization/depolymerization (effects on microtubule dynamics); binding to other proteins of the spindle; binding to proteins involved in cell-cycle control; serving as a substrate to other enzymes, such as kinases or proteases; and specific kinesin cellular activities such as spindle pole separation.

- [0108] Methods of performing motility assays are well known to those of skill in the art. [See e.g., Hall, et al. (1996), Biophys. J., 71: 3467-3476, Turner et al., 1996, Anal. Biochem. 242 (1):20-5; Gittes et al., 1996, Biophys. J. 70(l): 418-29; Shirakawa et al., 1995, J. Exp. Biol. 198: 1809-15; Winkelmann et al., 1995, Biophys. J. 68: 2444-53; Winkelmann et al., 1995, Biophys. J. 68: 72S.]
- [0109] Methods known in the art for determining ATPase hydrolysis activity also can be used. Solution based assays are particularly suitable (see, U.S. Patent No. 6,410,254); alternatively, conventional methods are used. For example,  $P_i$  release from kinesin can be quantified. In one embodiment, the ATPase hydrolysis activity assay utilizes 0.3 M PCA (perchloric acid) and malachite green reagent (8.27 mM sodium molybdate II, 0.33 mM malachite green oxalate, and 0.8 mM Triton X-100). To perform the assay, 10  $\mu$ L of reaction is quenched in 90  $\mu$ L of cold 0.3 M PCA. Phosphate standards are used so data can be converted to mM inorganic phosphate released. When all reactions and standards have been quenched in PCA, 100  $\mu$ L of malachite green reagent is added to the relevant wells in e.g., a microtiter plate. The mixture is developed for 10-15 minutes and the plate is read at an absorbance of 650 nm. When phosphate standards are used, absorbance readings can be converted to mM  $P_i$  and plotted over time. Additionally, ATPase assays known in the art include the luciferase assay.
- [0110] ATPase activity of kinesin motor domains also can be used to monitor the effects of modulating agents. In one embodiment ATPase assays of kinesin are performed in the absence of microtubules. In another embodiment, the ATPase assays are performed in the presence of microtubules. Different types of modulating agents can be detected in the above assays. In one particular embodiment, the effect of a modulating agent is independent of the concentration of microtubules and ATP. In another embodiment, the effect of the agents on kinesin ATPase can be decreased by increasing the concentrations of ATP, microtubules or both. In yet another embodiment, the effect of the modulating agent is increased by increasing concentrations of ATP, microtubules or both.
- [0111] Agents that modulate the biochemical activity of KSP *in vitro* may then be screened *in vivo*. Methods for testing such agents *in vivo* include assays of cell cycle distribution, cell viability, or the presence, morphology, activity, distribution or amount of mitotic spindles. Methods for monitoring cell cycle distribution of a cell population, for example, by flow cytometry, are well known to those skilled in the art, as are methods for determining cell viability. See, for example, WO 01/31335, entitled "Methods of Screening for Modulators of Cell Proliferation and Methods of Diagnosing Cell Proliferation States."

[0112] In addition to the assays described above, microscopic methods for monitoring spindle formation and malformation are well known to those of skill in the art (see, e.g., Whitehead and Rattner (1998), J. Cell Sci. 111:2551-61; Galgio et al, (1996) J. Cell Biol., 135:399-414).

[0113] The compounds of the invention inhibit KSP kinesin. One measure of inhibition,  $IC_{50}$ , is defined as the concentration of the compound at which the activity of KSP is decreased by fifty percent. Particularly suitable compounds have  $IC_{50}$ 's of less than about 1 mM, with more particularly suitable compounds having  $IC_{50}$ 's of less than about 100  $\mu$ M.  $IC_{50}$ 's of less than about 10 nM can be attained by certain compounds of the invention, and the pharmaceutically acceptable salts and solvates thereof, it being appreciated that a smaller  $IC_{50}$  is generally considered advantageous. Measurement of  $IC_{50}$  is done using an ATPase assay.

[0114] Another measure of inhibition is  $K_i$ . For compounds with  $IC_{50}$ 's less than 1  $\mu$ M, the  $K_i$  or  $K_d$  is defined as the dissociation rate constant for the interaction of the test compound with KSP. Particularly suitable compounds have  $K_i$ 's of less than about 100  $\mu$ M, more particularly suitable compounds having  $K_i$ 's of less than about 10  $\mu$ M.  $K_i$ 's of less than about 10 nM can be attained by certain compounds of the invention, and the pharmaceutically acceptable salts and solvates thereof, it being appreciated that a smaller  $K_i$  is generally considered advantageous. The  $K_i$  for a compound is determined from the  $IC_{50}$  based on three assumptions. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[ 1 - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4 E_0 I_0}}{2 E_0} \right]$$

Where  $V$  is the observed rate,  $V_{\max}$  is the rate of the free enzyme,  $I_0$  is the inhibitor concentration,  $E_0$  is the enzyme concentration, and  $K_d$  is the dissociation constant of the enzyme-inhibitor complex.

[0115] Another measure of inhibition is  $GI_{50}$ , defined as the concentration of the compound that results in a decrease in the rate of cell growth by fifty percent. Anti-proliferative compounds that have been successfully applied in the clinic to treatment of cancer (cancer chemotherapeutics) have  $GI_{50}$ 's that vary greatly. For example, in A549 cells, paclitaxel  $GI_{50}$  is

4 nM, doxorubicin is 63 nM, 5-fluorouracil is 1  $\mu$ M, and hydroxyurea is 500  $\mu$ M (data provided by National Cancer Institute, Developmental Therapeutic Program, <http://dtp.nci.nih.gov/>). Therefore, compounds that inhibit cellular proliferation at virtually any concentration may be useful. Particularly suitable compounds have  $GI_{50}$ 's of less than about 1 mM, with more particularly suitable compounds having a  $GI_{50}$  of less than about 10  $\mu$ M.  $GI_{50}$ 's of less than about 10 nM can be attained by certain compounds of the invention, and the pharmaceutically acceptable salts and solvates thereof, it being appreciated that a smaller  $GI_{50}$  is generally considered advantageous. Measurement of  $GI_{50}$  is done using a cell proliferation assay.

[0116] Testing for growth inhibition using cell lines (such as MCF-7/ADR-RES and HCT1 5) that express P-glycoprotein (also known as Multi-drug Resistance, or MDR<sup>+</sup>), which conveys resistance to other chemotherapeutic drugs, such as paclitaxel, can identify anti-mitotic agents that inhibit cell proliferation and are not subject to resistance by overexpression of MDR<sup>+</sup> by drug-resistant tumor lines.

[0117] *In vitro* potency of small molecule inhibitors is determined by assaying human ovarian cancer cells (SKOV3) for viability following a 72-hour exposure to a 9-point dilution series of compound. Cell viability is determined by measuring the absorbance of formazon, a product formed by the bioreduction of MTS/PMS, a commercially available reagent. Each point on the dose-response curve is calculated as a percent of untreated control cells at 72 hours minus background absorption (complete cell kill).

[0118] To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound or composition of the invention is added to the assay. Alternatively, a composition of a compound of the invention bound to a solid support can be made, and KSP added to the assay. Classes of compounds among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[0119] The determination of the binding of the mitotic agent to KSP may be done in a number of ways. In a particular embodiment, the compound of the invention is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this may be done by attaching all or a portion of KSP to a solid support, adding a labeled compound (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the

amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

[0120] By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[0121] In some embodiments, only one of the components is labeled. For example, the kinesin proteins may be labeled at tyrosine positions using  $^{125}\text{I}$ , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using  $^{125}\text{I}$  for the proteins, for example, and a fluorophor for the anti-mitotic agents.

[0122] The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. Screens of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, particular embodiments exclude molecules already known to bind to that protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Particular embodiments of assays herein include candidate agents that do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous" agents. In another particular embodiment, exogenous agents further exclude antibodies to KSP.

[0123] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, particularly small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, especially at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures

substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

[0124] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

[0125] Competitive screening assays can be done by combining KSP and a drug candidate in a first sample. A second sample may be made combining a compound of the invention, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change or difference in binding between the two samples indicates the presence of an agent capable of binding to KSP and potentially modulating its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

[0126] In a particularly suitable embodiment, the binding of the candidate agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[0127] In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if present. Incubations can be performed at any temperature that facilitates optimal activity, typically between 4 and 40°C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[0128] In a particularly suitable embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is



binding to KSP and thus is capable of binding to, and potentially modulating, the activity of KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[0129] In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to KSP.

[0130] It may be of value to identify the binding site of KSP. This can be done in a variety of ways. In one embodiment, once KSP has been identified as binding to the compound, KSP is fragmented or modified and the assays repeated to identify the necessary components for binding.

[0131] Modulation is tested by screening for candidate agents capable of modulating the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both *in vitro* screening methods and *in vivo* screening of cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

[0132] Alternatively, differential screening may be used to identify drug candidates that bind to the native KSP, but cannot bind to modified KSP.

[0133] Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[0134] A variety of other reagents can be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors,

nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

#### Formulation and Administration

- [0135] The compounds, pharmaceutically acceptable salts and solvates of Formula I and II are administered at a therapeutically effective dosage, e.g., a dosage sufficient to provide treatment for the disease states previously described. Human dosage levels are typically determined by escalating dose ranging studies conducted in accordance with current Good Clinical Practice, FDA and local guidelines. The amount of active compound administered will, of course, be dependent on the subject and disease state being treated, the severity of the affliction, the manner and schedule of administration and the judgment of the prescribing physician.
- [0136] The administration of the compounds and pharmaceutical formulations of the present invention can be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the compound or composition may be directly applied as a solution or spray.
- [0137] Pharmaceutical formulations include a compound of Formula I or II or a pharmaceutically acceptable salt or solvate thereof, and one or more pharmaceutically acceptable excipients. As is known in the art, pharmaceutical excipients are secondary ingredients that function to enable or enhance the delivery of a drug or medicine in a variety of dosage forms (e.g.: oral forms such as tablets, capsules, and liquids; topical forms such as dermal, ophthalmic, and otic forms; suppositories; injectables; respiratory forms and the like). Pharmaceutical excipients include inert or inactive ingredients, synergists or chemicals that substantively contribute to the medicinal effects of the active ingredient. For example, pharmaceutical excipients may function to improve flow characteristics, product uniformity, stability, taste, or appearance, to ease handling and administration of dose, for convenience of use, or to control bioavailability. While pharmaceutical excipients are commonly described as being inert or inactive, it is appreciated in the art that there is a relationship between the properties of the pharmaceutical excipients and the dosage forms containing them.
- [0138] Pharmaceutical excipients suitable for use as carriers or diluents are well known in the art, and may be used in a variety of formulations. See, e.g., Remington's Pharmaceutical Sciences, 18th Edition, A. R. Gennaro, Editor, Mack Publishing Company (1990); Remington: The Science and Practice of Pharmacy, 20th Edition, A. R. Gennaro, Editor, Lippincott Williams & Wilkins (2000); Handbook of Pharmaceutical Excipients, 3rd Edition, A. H. Kibbe,

Editor, American Pharmaceutical Association, and Pharmaceutical Press (2000); and Handbook of Pharmaceutical Additives, compiled by Michael and Irene Ash, Gower (1995). The concentration of a therapeutically active agent in a formulation can vary widely, from about 0.1 to 99.9 wt.%, depending on the nature of the formulation.

[0139] Oral solid dosage forms such as tablets will typically comprise one or more pharmaceutical excipients, which may for example help impart satisfactory processing and compression characteristics, or provide additional desirable physical characteristics to the tablet. Such pharmaceutical excipients may be selected from diluents, binders, glidants, lubricants, disintegrants, colorants, flavorants, sweetening agents, polymers, waxes or other solubility-modulating materials.

[0140] Dosage forms for parenteral administration will generally comprise fluids, particularly intravenous fluids, i.e., sterile solutions of simple chemicals such as sugars, amino acids or electrolytes, which can be easily carried by the circulatory system and assimilated. Such fluids are typically prepared with water for injection USP. Fluids used commonly for intravenous (IV) use are disclosed in Remington, The Science and Practice of Pharmacy [full citation previously provided], and include:

- alcohol, e.g., 5% alcohol (e.g., in dextrose and water ("D/W") or D/W in normal saline solution ("NSS"), including in 5% dextrose and water ("D5/W"), or D5/W in NSS);
- synthetic amino acid such as Aminosyn, FreAmine, Travasol, e.g., 3.5 or 7; 8.5; 3.5, 5.5 or 8.5 % respectively;
- ammonium chloride e.g., 2.14%;
- dextran 40, in NSS e.g., 10% or in D5/W e.g., 10%;
- dextran 70, in NSS e.g., 6% or in D5/W e.g., 6%;
- dextrose (glucose, D5/W) e.g., 2.5-50%;
- dextrose and sodium chloride e.g., 5-20% dextrose and 0.22-0.9% NaCl;
- lactated Ringer's (Hartmann's) e.g., NaCl 0.6%, KCl 0.03%,  $\text{CaCl}_2$  0.02%;
- lactate 0.3%;
- mannitol e.g., 5%, optionally in combination with dextrose e.g., 10% or NaCl e.g., 15 or 20%;
- multiple electrolyte solutions with varying combinations of electrolytes, dextrose, fructose, invert sugar Ringer's e.g., NaCl 0.86%, KCl 0.03%,  $\text{CaCl}_2$  0.033%;
- sodium bicarbonate e.g., 5%;
- sodium chloride e.g., 0.45, 0.9, 3, or 5%;

- sodium lactate e.g., 1/6 M; and
- sterile water for injection

The pH of such IV fluids may vary, and will typically be from 3.5 to 8 as known in the art.

[0141] The compounds, pharmaceutically acceptable salts and solvates of the invention can be administered alone or in combination with other treatments, i.e., radiation, or other therapeutic agents, such as the taxane class of agents that appear to act on microtubule formation or the camptothecin class of topoisomerase I inhibitors. When so-used, other therapeutic agents can be administered before, concurrently (whether in separate dosage forms or in a combined dosage form), or after administration of an active agent of the present invention.

[0142] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes.

## EXAMPLES

### Example 1

#### 3-Benzyl-7-chloro-1-pyrrolidine-2-yl-2,3-dihydro-1H-quinazoline-4-one

[0143] 1A. Formula 103 where T is a covalent bond; W, X, Y and Z are -C=; U'-V' is -N(R<sup>6'</sup>)-CR<sup>6'</sup>R<sup>1'</sup>; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, and R<sup>5</sup> to R<sup>1'</sup> are H; R<sup>3</sup> is chloro; and R<sup>6'</sup> is Boc A solution of pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (6.0 g, 28 mmol), 2-amino-4-chloro-benzoic acid methyl ester (4.6 g, 25 mmol), PyBroP (14.4 g, 31 mmol), diisopropylethylamine (6 mL) in pyridine (60 mL) was stirred overnight at room temperature, after which HPLC/MS showed complete conversion of starting material. The solvents were removed under vacuum and the residue partitioned between ethyl acetate (100 mL) and water (50 mL). The organic layer was washed with 1M HCl (50 mL) and saturated NaHCO<sub>3</sub> (50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated under vacuum. Silica gel chromatography using 30% ethyl acetate/hexanes as eluent gave 9.3 g (87% yield) of the desired product of Formula 103, 2-(5-chloro-2-methoxycarbonyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester, as a yellow foam. LRMS (MH<sup>+</sup>) *m/z* 383.

[0144] 1B. Formula 104 where T is a covalent bond; W, X, Y and Z are  $-C=$ ; U'-V' is  $-N(R^6)-CR^6R^7$ ;  $R^1, R^2, R^4$ , and  $R^a$  to  $R^f$  are H;  $R^3$  is chloro; and  $R^6$  is Boc: A solution of the pure compound of Formula 103 (9.3 g, 24 mmol) dissolved in 125 mL of ethanol was treated with NaOH solution (3.1g in 16 mL of  $H_2O$ ). The mixture was stirred at room temperature overnight and then concentrated under vacuum. The residue was treated with a solution of saturated NaCl (200 mL) and concentrated phosphoric acid (5 mL) for a few minutes and then extracted with dichloromethane (2 x 200 mL). The organic fractions were combined, dried over  $MgSO_4$ , filtered and concentrated give 8.7 g (97% yield) of the clean desired product of Formula 104, 2-(5-chloro-2-carboxy-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester, as an orange foam. LRMS (M)  $m/z$  368.

[0145] 1C. Formula 105 where T is a covalent bond; W, X, Y and Z are  $-C=$ ; U'-V' is  $-N(R^6)-CR^6R^7$ ;  $R^1, R^2, R^4$ , and  $R^a$  to  $R^f$  are H;  $R^3$  is chloro;  $R^5$  is benzyl, and  $R^6$  is Boc:

To a solution of the compound of Formula 104 (8.7 g, 24 mmol) in dichloromethane (250 mL) was added EDC (14.85 g, 77 mmol), and diisopropylethylamine (14.1 mL, 81 mmol). After stirring at room temperature for 1.5 hours, benzylamine (8.4 mL, 77 mmol) was added and the reaction stirred for an additional 48 hours. The mixture was washed with water (2 x 50 mL), 10% phosphoric acid (2 x 50 mL), and saturated  $NaHCO_3$  (50 mL). The organic phase was dried over  $MgSO_4$ , filtered and concentrated under vacuum to give a mixture of 9.4 g (88% yield) of the desired product of Formula 105, 2-(2-benzylcarbamoyl-5-chloro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester, as a light yellow foam, together with a small amount of 2-(3-benzyl-7-chloro-4-oxo-3,4-dihydro-2H-quinzoline-1-yl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (Formula 106), which was taken on without further purification. LRMS ( $MH^+$ )  $m/z$  458.

[0146] 1D. Formula 106 where T is a covalent bond; W, X, Y and Z are  $-C=$ ; U'-V' is  $-N(R^6)-CR^6R^7$ ;  $R^1, R^2, R^4$ , and  $R^a$  to  $R^f$  are H;  $R^3$  is chloro;  $R^5$  is benzyl, and  $R^6$  is Boc:

The mixture of Formulae 105 and 106 (5.4 g, 12 mmol) was stirred in ethylene glycol (32 mL) with  $K_2CO_3$  (27 g) at 120°C overnight, after which the solution was allowed to cool to room temperature and was extracted with dichloromethane (3 x 200 mL). The organic fractions were combined, dried over  $MgSO_4$ , filtered and concentrated under vacuum. The resulting residue was purified by silica gel chromatography using 20% ethyl acetate/hexanes as eluent to give 1.4 g (27% yield) of the pure desired product of Formula 106, 2-(3-benzyl-7-chloro-4-oxo-3,4-dihydro-2H-quinzoline-1-yl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester. LRMS ( $MH^+$ )  $m/z$  439.

- [0147] 1E. Formula I where U-V is -N(R<sup>6</sup>)-CR<sup>6</sup>R<sup>1</sup>; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>6</sup> and R<sup>a</sup> to R<sup>f</sup> are H; R<sup>3</sup> is chloro; and R<sup>5</sup> is benzyl: The pure compound of Formula 106 (446 mg, 1.0 mmol) was treated with 20 mL of a 1:1 mixture of TFA/dichloromethane for 1 hour at room temperature. The solution was concentrated under vacuum and partitioned between dichloromethane (20 mL) and saturated NaHCO<sub>3</sub> (10 mL). The aqueous layer was extracted with dichloromethane (10 mL) and the combined organic layers were dried over K<sub>2</sub>CO<sub>3</sub>, filtered and concentrated to give 344 mg (100% yield) of the desired product of Formula I, 3-benzyl-7-chloro-1-pyrrolidine-2-yl-2,3-dihydro-1*H*-quinazoline-4-one, which was taken on without further purification. LRMS (MH<sup>+</sup>) *m/z* 340.

### Example 2

#### 3-Benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one

- [0148] Formula I where U-V is -N(R<sup>6</sup>)-CR<sup>6</sup>R<sup>1</sup>; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, and R<sup>a</sup> to R<sup>f</sup> are H; R<sup>3</sup> is chloro; R<sup>5</sup> is benzyl; and R<sup>6</sup> is *p*-methyl-benzoyl: To a solution of crude 3-benzyl-7-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1*H*-quinazoline-4-one (250 mg, 0.73 mmol) in dichloromethane (4.5 mL) was added *p*-toluoyl chloride (0.29 mL, 2.2 mmol) and diisopropylethylamine (0.77 mL, 4.4 mmol). The mixture was stirred for 1 hour and then partitioned between saturated NaHCO<sub>3</sub> (10 mL) and ethyl acetate (20 mL). The organic phase was washed with saturated NaHCO<sub>3</sub> (3 x 4 mL) and saturated NaCl (2 x 4 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude residue was purified by preparative reverse phase C18 HPLC (10-98% acetonitrile/water) to give 223 mg (66% yield) of the pure desired product of Formula I, 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one. LRMS (MH<sup>+</sup>) *m/z* 459.

### Example 3

#### 3-Benzyl-7-chloro-2-[1-(4-methyl-benzyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one

- [0149] Formula I where U-V is -N(R<sup>6</sup>)-CR<sup>6</sup>R<sup>1</sup>; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, and R<sup>a</sup> to R<sup>f</sup> are H; R<sup>3</sup> is chloro; R<sup>5</sup> is benzyl; and R<sup>6</sup> is *p*-methyl-benzyl: Crude 3-benzyl-7-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1*H*-quinazoline-4-one (170 mg, 0.50 mmol) dissolved in dichloromethane (5 mL) was treated with *p*-tolualdehyde (80  $\mu$ L, 0.7 mmol) and NaHB(OAc)<sub>3</sub> (350 mg, 1.65 mmol), and the mixture was stirred for 40 minutes at room temperature. The reaction was

quenched with saturated  $\text{NaHCO}_3$  (2.2 mL) and the aqueous phase extracted with dichloromethane (25 mL). The organic layers were combined and dried over  $\text{Na}_2\text{SO}_3$ , filtered and concentrated under vacuum. The crude residue was purified by silica gel chromatography using a stepwise gradient of 20% ethyl acetate/hexanes to 50% methanol/ethyl acetate as eluent to give 52.5 mg (24 % yield) of desired product of Formula I, 3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-pyrrolidin-2-yl]-3H-quinazolin-4-one. LRMS ( $\text{MH}^+$ )  $m/z$  445.

### Example 4

#### Other Compounds of Formulae I and II

[0150] 4A. Formula I where U-V is  $-\text{CR}^{\text{e}}\text{R}^{\text{f}}-\text{CR}^{\text{g}}\text{R}^{\text{h}}-\text{N}(\text{R}^{\text{b}})-$ ;  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^{\text{b}}$  and  $\text{R}^{\text{c}}$  to  $\text{R}^{\text{h}}$  are H;  $\text{R}^3$  is chloro;  $\text{R}^5$  is methyl; and  $\text{R}^{\text{a}}$  is phenethyl; and  $\text{R}^6$  is p-methyl-benzoyl By following the procedure described in Example 1 and substituting pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester with 3-phenethyl-piperidine-1,4-dicarboxylic acid *tert*-butyl ester (in Example 1A), and substituting benzylamine with methylamine (in Example 1C) there is obtained: 7-chloro-2-[3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one.

[0151] 4B. Formula I where U-V is  $-\text{CR}^{\text{e}}\text{R}^{\text{f}}-\text{CR}^{\text{g}}\text{R}^{\text{h}}-\text{N}(\text{R}^{\text{b}})-$ ;  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^{\text{b}}$  and  $\text{R}^{\text{c}}$  to  $\text{R}^{\text{h}}$  are H;  $\text{R}^3$  is chloro;  $\text{R}^5$  is methyl; and  $\text{R}^{\text{a}}$  is phenethyl; and  $\text{R}^6$  is p-methyl-benzoyl: By following the procedure described in Example 2 and substituting 3-benzyl-7-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1H-quinazoline-4-one with 7-chloro-2-[3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one, there is obtained: 7-chloro-2-[1-(4-methyl-benzoyl)-3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one.

[0152] 4C. Formula I where U-V is  $-\text{CR}^{\text{e}}\text{R}^{\text{f}}-\text{CR}^{\text{g}}\text{R}^{\text{h}}-\text{N}(\text{R}^{\text{b}})-$ ;  $\text{R}^1, \text{R}^4, \text{R}^{\text{b}}$  and  $\text{R}^{\text{c}}$  to  $\text{R}^{\text{h}}$  are H;  $\text{R}^2$  and  $\text{R}^3$  are methoxy;  $\text{R}^5$  is methyl; and  $\text{R}^{\text{a}}$  is phenethyl; and  $\text{R}^6$  is benzyl: By following the procedure described in Example 3, substituting 3-benzyl-7-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1H-quinazoline-4-one with 6,7-dimethoxy-2-[3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one, and substituting *p*-tolualdehyde with benzaldehyde, there is obtained 6,7-dimethoxy-2-[1-benzyl-3-phenethyl-piperidin-4-yl]-3-methyl-3H-quinazolin-4-one.

[0153] 4D. Compounds of Formula II Varying T By following the procedure described in Example 1 and substituting pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester with the following:

- 2-carboxymethyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester;
- 2-(2-carboxy-acetyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester; and
- 2-(2-carboxy-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester,

there are obtained the following respective compounds:

- 3-benzyl-7-chloro-2-pyrrolidine-2-ylmethyl-3*H*-quinazoline-4-one;
- 3-benzyl-7-chloro-2-(2-oxo-2-pyrrolidine-2-yl-ethyl)-3*H*-quinazoline-4-one; and
- 3-benzyl-7-chloro-2-piperidine-2-ylethyl-3*H*-quinazoline-4-one.

[0154] 4E. Compounds of Formula II Varying W, X, Y and Z By following the procedure described in Example 1 and substituting 2-amino-4-chloro-benzoic acid methyl ester with the following:

- 4-amino-6-chloro-nicotinic acid methyl ester;
- 3-amino-pyrazine-2-carboxylic acid;
- 3-amino-1,4-dihydro-pyridine-2-carboxylic acid;
- 2-amino-cyclopent-1-enecarboxylic acid;
- 4-amino-2,5-dihydro-furan-3-carboxylic acid; and
- 3-amino-1*H*-pyrrole-2-carboxylic acid,

there are obtained the following respective compounds:

- 3-benzyl-7-chloro-1-pyrrolidine-2-yl-2,3-dihydro-1*H*-pyrido[4,3-*d*]pyrimidin-4-one;
- 3-benzyl-1-pyrrolidin-2-yl-1*H*-pteridin-4-one;
- 3-benzyl-1-pyrrolidin-2-yl-2,3,5,6,7,8-hexahydro-1*H*-pyrido[3,2-*d*]pyrimidin-4-one;
- 3-benzyl-1-pyrrolidin-2-yl-1,2,3,5,6,7-hexahydro-cyclopentapyrimidin-4-one;
- 3-benzyl-1-pyrrolidin-2-yl-2,3,5,7-tetrahydro-1*H*-furo[3,4-*d*]pyrimidin-4-one; and
- 3-benzyl-1-pyrrolidin-2-yl-1,2,3,5-tetrahydro-pyrrolo[3,2-*d*]pyrimidin-4-one.

### Example 5

#### 3-Benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one

[0155] 5A. Formula 202 where T is a covalent bond; U-V is -N(R<sup>6</sup>)-CR<sup>6</sup>R<sup>1</sup>-CR<sup>9</sup>R<sup>h</sup>-; R<sup>a</sup> to R<sup>h</sup> are H; R<sup>3</sup> is chloro; and R<sup>8</sup> is *p*-methyl-benzoyl: To a solution of piperidine-2-carboxylic acid (3.5 g, 15 mmol) in 2M NaOH (70 mL) at 0°C, was added *p*-toluoyl chloride (2.2 mL, 17 mmol) in 10 equal portions over 1 hour. The mixture was then stirred an additional 1 hour at room temperature. The reaction was cooled to 0°C and partitioned between 1N HCl and dichloromethane. The organic layer was dried over MgSO<sub>4</sub>, filtered and concentrated under vacuum to give 4.7 g of oil. 2.0 g of this material was recrystallized from 95% ethanol (2.5 mL) to give 0.86 g of the desired pure crystalline product of Formula 202, 1-(4-methyl-benzoyl)-piperidine-2-carboxylic acid. LRMS (MH<sup>+</sup>) *m/z* 248.



[0156] 5B. Formula 203 where T is a covalent bond; W, X, Y and Z are -C=; U-V is -N(R<sup>6</sup>)-CR<sup>a</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup> and R<sup>a</sup> to R<sup>h</sup> are H; R<sup>3</sup> is chloro; and R<sup>6</sup> is p-methyl-benzoyl:

In an ice bath at -15°C, pure compound of Formula 202 (7.2 g, 29 mmol) was combined with 2-amino-4-chloro-benzoic acid methyl ester (4.5 g, 24 mmol), PyBroP (2.3 g, 4.9 mmol) and diisopropylethylamine (0.95 mL, 5.4 mmol) in DMF (30 mL). The resulting mixture was stirred overnight, allowing the temperature to rise to room temperature with the bath. The solution was concentrated under vacuum and the residue partitioned between ethyl acetate (200 mL) and water (200 mL). The organic layer was washed with water (3 x 100 mL) and concentrated under vacuum. The resulting residue was purified by silica gel chromatography using 40% ethyl acetate/hexanes as eluent to give 2.66 g (21% yield) of the desired product of Formula 203, 4-chloro-2-([1-(4-methyl-benzoyl)-piperidine-2-carbonyl]amino)-benzoic acid methyl ester, as a crystalline solid. LRMS (MH<sup>+</sup>) m/z 472.

[0157] 5C. Formula 205 where T is a covalent bond; W, X, Y and Z are -C=; U-V is -N(R<sup>6</sup>)-CR<sup>a</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup> and R<sup>a</sup> to R<sup>h</sup> are H; R<sup>3</sup> is chloro; R<sup>5</sup> is benzyl; and R<sup>6</sup> is p-methyl-benzoyl:

A mixture of the compound of Formula 203 (2.66 g, 6.7 mmol), 1M NaOH (27 mL), methanol (54 mL), and dioxane (36 mL) was stirred at room temperature for 7 hours. The solvents were evaporated and the residue portioned between dichloromethane (75 mL), saturated NaCl (50 mL), and concentrated HCl (3 mL). The layers were separated and the organic layer dried over MgSO<sub>4</sub> and concentrated under vacuum to give the corresponding compound of Formula 204, 4-chloro-2-([1-(4-methyl-benzoyl)-piperidine-2-carbonyl]amino)-benzoic acid, which was carried forward without further purification. This crude product was redissolved in dichloromethane and treated with EDC (3.84 g, 20 mmol), and diisopropylethylamine (3.50 mL, 20 mmol). After stirring 20 minutes, benzylamine (2.20 mL, 20 mmol) was added and the mixture stirred overnight at room temperature. The mixture was washed with water (2 x 50 mL) and 1M HCl, dried over MgSO<sub>4</sub>, filtered and concentrated under vacuum to give 1.5 g (47% yield) of the desired product of Formula 205, 1-(4-methyl-benzoyl)-piperidine-2-carboxylic acid (2-benzylcarbonyl-5-chloro-phenyl-amide, a crude product that was found to be 80% pure by reverse phase HPLC. The product was taken on without further purification. LRMS (MH<sup>+</sup>) m/z 491.

[0158] 5D. Formula I where U-V is -N(R<sup>6</sup>)-CR<sup>a</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-; R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup> and R<sup>a</sup> to R<sup>h</sup> are H; R<sup>3</sup> is chloro; R<sup>5</sup> is benzyl; and R<sup>6</sup> is p-methyl-benzoyl:

A mixture of the crude product of Formula 205 (170 mg, 0.35 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.8 g, 5.8 mmol) in ethylene glycol (4 mL) was stirred at 120°C overnight. After cooling to room temperature, the solution was diluted with

water (30 mL), and extracted with dichloromethane (30 mL) and ethyl acetate (30 mL). The organic layers were combined and washed with water (2 x 5 mL) and saturated NaCl (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum to give a turbid yellow oil. The crude material was purified by reverse phase C18 HPLC (10-98% acetonitrile/water) to give 19 mg (12% yield) of the desired product of Formula I, 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one. LRMS (MH<sup>+</sup>) *m/z* 473.

### Example 6

#### Other Compounds of Formulae I and II

[0159] 6A. Formula I Varying R<sup>6</sup>: By following the procedure described in Example 5 and substituting *p*-toluoyl chloride with the following:

- benzyl chloride;
- benzoyl chloride; and
- *p*-tolyl chloride,

there are obtained the following respective compounds:

- 3-benzyl-7-chloro-2-(1-benzyl-piperidin-2-yl)-3*H*-quinazolin-4-one;
- 3-benzyl-7-chloro-2-(1-benzoyl-piperidin-2-yl)-3*H*-quinazolin-4-one; and
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-piperidin-2-yl]-3*H*-quinazolin-4-one.

[0160] 6B. Compounds of Formula II Varying T By following the procedure described in Example 5 and substituting piperidine-1,2-dicarboxylic acid with the following:

- 2-carboxymethyl-pyrrolidine-1-carboxylic acid;
- 2-(2-carboxy-acetyl)-pyrrolidine-1-carboxylic acid; and
- 2-(2-carboxy-ethyl)-piperidine-1-carboxylic acid,

there are obtained the following respective compounds:

- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidine-2-ylmethyl]-3*H*-quinazoline-4-one;
- 3-benzyl-7-chloro-2-[2-[1-(4-methyl-benzoyl)-pyrrolidine-2-yl]-2-oxo-ethyl]-3*H*-quinazoline-4-one; and
- 3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidine-2-ylethyl]-3*H*-quinazoline-4-one.

[0161] 6C. Compounds of Formula II Varying W, X, Y and Z By following the procedure described in Example 5 and substituting 2-amino-4-chloro-benzoic acid methyl ester with the following:

- 4-amino-6-chloro-nicotinic acid methyl ester;
- 3-amino-pyrazine-2-carboxylic acid;

- 3-amino-1,4-dihydro-pyridine-2-carboxylic acid;
- 2-amino-cyclopent-1-enecarboxylic acid;
- 4-amino-2,5-dihydro-furan-3-carboxylic acid; and
- 3-amino-1*H*-pyrrole-2-carboxylic acid,

there are obtained the following respective compounds:

- 3-benzyl-7-chloro-1-(4-methyl-benzoyl)-pyrrolidine-2-yl-2,3-dihydro-1*H*-pyrido[4,3-*d*]pyrimidin-4-one;
- 3-benzyl-1-(4-methyl-benzoyl)-pyrrolidin-2-yl-1*H*-pteridin-4-one;
- 3-benzyl-1-(4-methyl-benzoyl)-pyrrolidin-2-yl-2,3,5,6,7,8-hexahydro-1*H*-pyrido[3,2-*d*]pyrimidin-4-one;
- 3-benzyl-1-(4-methyl-benzoyl)-pyrrolidin-2-yl-1,2,3,5,6,7-hexahydro-cyclopentapyrimidin-4-one;
- 3-benzyl-1-(4-methyl-benzoyl)-pyrrolidin-2-yl-2,3,5,7-tetrahydro-1*H*-furo[3,4-*d*]pyrimidin-4-one; and
- 3-benzyl-1-(4-methyl-benzoyl)-pyrrolidin-2-yl-1,2,3,5-tetrahydro-pyrrolo[3,2-*d*]pyrimidin-4-one.

### Example 7

#### Induction of Mitotic Arrest in Cell Populations Treated with a KSP Inhibitor

[0162] FACS analysis to determine cell cycle stage by measuring DNA content is performed as follows. Skov-3 cells (human ovarian cancer) are split 1:10 for plating in 10cm dishes and grown to subconfluence with RPMI 1640 medium containing 5% fetal bovine serum (FBS). The cells are then treated with either 10nM paclitaxel, 400nM test compound, 200nM test compound, or 0.25% DMSO (vehicle for compounds) for 24 hours. A well known anti-mitotic agent, such as paclitaxel, is used as a positive control. Cells are then rinsed off the plates with PBS containing 5mM EDTA, pelleted, washed once in PBS containing 1% FCS, and then fixed overnight in 85% ethanol at 4°C. Before analysis, the cells are pelleted, washed once, and stained in a solution of 10µg propidium iodide and 250µg of ribonuclease (RNase) A per milliliter at 37°C for half an hour. Flow cytometry analysis is performed on a Becton-Dickinson FACScan, and data from 10,000 cells per sample is analyzed with Modfit software.

#### Monopolar Spindle Formation following Application of a Quinazolinone KSP Inhibitor

[0163] To determine the nature of G2/M accumulation, human tumor cell lines Skov-3

(ovarian), HeLa (cervical), and A549 (lung) are plated in 96-well plates at densities of 4,000 cells per well (SKOV-3 & HeLa) or 8,000 cells per well (A549), allowed to adhere for 24 hours, and treated with various concentrations of the test compounds for 24 hours. Cells are fixed in 4% formaldehyde and stained with antitubulin antibodies (subsequently recognized using fluorescently-labeled secondary antibody) and Hoechst dye (which stains DNA). The cells can be visually inspected to assess the effects of the test compounds. For example, microinjection of anti-KSP antibodies causes mitotic arrest with arrested cells displaying monopolar spindles.

### Example 8

#### Inhibition of Cellular Proliferation in Tumor Cell Lines Treated with KSP Inhibitors.

[0164] Cells are plated in 96-well plates at densities from 1000-2500 cells/well (depending on the cell line) and allowed to adhere/grow for 24 hours. They are then treated with various concentrations of test compound for 48 hours. The time at which compounds are added is considered  $T_0$ . A tetrazolium-based assay using the reagent 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) (U.S. Patent No. 5,185,450) (see Promega product catalog #G3580, CellTiter 96® AQueous One Solution Cell Proliferation Assay) is used to determine the number of viable cells at  $T_0$  and the number of cells remaining after 48 hours compound exposure. The number of cells remaining after 48 hours is compared to the number of viable cells at the time of test compound addition, allowing for calculation of growth inhibition. The growth over 48 hours of cells in control wells treated with vehicle only (0.25% DMSO) is considered 100% growth and the growth of cells in wells with compounds is compared to this. Active KSP inhibitors inhibit cell proliferation in one or more human tumor cell lines of the following tumor types: lung (NCI-H460, A549), breast (MDA-MB-231, MCF-7, MCF-7/ADR-RES), colon (HT29, HCT15), ovarian (SKOV-3, OVCAR-3), leukemia (HL-60(TB), K-562), central nervous system (SF-268), renal (A498), osteosarcoma (U2-OS), and cervical (HeLa), and mouse tumor line (B16, melanoma).

[0165] Calculation Of  $GI_{50}$ : A  $GI_{50}$  is calculated by plotting the concentration of compound in  $\mu M$  vs the percentage of cell growth of cell growth in treated wells. The  $GI_{50}$  calculated for the compounds is the estimated concentration at which growth is inhibited by 50% compared to control, i.e., the concentration at which:

$$100 \times [(Treated_{48} - T_0) / (Control_{48} - T_0)] = 50.$$

All concentrations of compounds are tested in duplicate and controls are averaged over 12 wells. A very similar 96-well plate layout and  $GI_{50}$  calculation scheme is used by the National Cancer Institute (see Monks, et al., J. Natl. Cancer Inst. 83:757-766 (1991)). However, the method by which the National Cancer Institute quantitates cell number does not use MTS, but instead employs alternative methods.

[0166] Calculation Of  $IC_{50}$ : Measurement of a compound's  $IC_{50}$  for KSP activity uses an ATPase assay. The following solutions are used: Solution 1 consists of 3 mM phosphoenolpyruvate potassium salt (Sigma P-7127), 2 mM ATP (Sigma A-3377), 1 mM IDTT (Sigma D-9779), 5  $\mu$ M paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM  $MgCl_2$  (VWR JT400301), and 1 mM EGTA (Sigma E3889). Solution 2 consists of 1 mM NADH (Sigma N8129), 0.2 mg/ml BSA (Sigma A7906), pyruvate kinase 7U/ml, L-lactate dehydrogenase 10 U/ml (Sigma P0294), 100 nM KSP motor domain, 50  $\mu$ g/ml microtubules, 1 mM DTT (Sigma D9779), 5  $\mu$ M paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM  $MgCl_2$  (VWR JT4003-01), and 1 mM EGTA (Sigma E3889). Serial dilutions (8-12 two-fold dilutions) of the composition are made in a 96-well microtiter plate (Corning Costar 3695) using Solution 1. Following serial dilution each well has 50  $\mu$ l of Solution 1. The reaction is started by adding 50  $\mu$ l of Solution 2 to each well. This can be done with a multichannel pipettor either manually or with automated liquid handling devices. The microtiter plate is then transferred to a microplate absorbance reader and multiple absorbance readings at 340 nm are taken for each well in a kinetic mode. The observed rate of change, which is proportional to the ATPase rate, is then plotted as a function of the compound concentration. For a standard  $IC_{50}$  determination the data acquired is fit by the following four parameter equation using a nonlinear fitting program (e.g., Grafit 4):

$$y = \frac{\text{Range}}{1 + \left( \frac{x}{IC_{50}} \right)^s} + \text{Background}$$

where y is the observed rate and x the compound concentration.

**Example 9****Inhibition of Cellular Viability in Tumor Cell Lines Treated with KSP Inhibitors.****[0167]      Materials and Solutions:**

- Cells: SKOV3, Ovarian Cancer (human).
- Media: Phenol Red Free RPMI + 5% Fetal Bovine Serum + 2mM L-glutamine.
- Colorimetric Agent for Determining Cell Viability: Promega MTS tetrazolium compound.
- Control Compound for max cell kill: Topotecan, 1 $\mu$ M.

**[0168]      Procedure: Day 1 - Cell Plating:**      Adherent SKOV3 cells are washed with 10mLs of PBS followed by the addition of 2mLs of 0.25% trypsin and incubation for 5 minutes at 37°C. The cells are rinsed from the flask using 8 mL of media (phenol red-free RPMI+ 5%FBS) and transferred to fresh flask. Cell concentration is determined using a Coulter counter and the appropriate volume of cells to achieve 1000 cells/100 $\mu$ L is calculated. 100  $\mu$ L of media cell suspension (adjusted to 1000 cells/100  $\mu$ L) is added to all wells of 96-well plates, followed by incubation for 18 to 24 hours at 37°C, 100% humidity, and 5% CO<sub>2</sub>, allowing the cells to adhere to the plates.

**[0169]      Procedure: Day 2 – Compound Addition:**      To one column of the wells of an autoclaved assay block are added an initial 2.5  $\mu$ L of test compound(s) at 400X the highest desired concentration. 1.25  $\mu$ L of 400X (400 $\mu$ M) Topotecan is added to other wells (ODs from these wells are used to subtract out for background absorbance of dead cells and vehicle). 500  $\mu$ L of media without DMSO are added to the wells containing test compound, and 250  $\mu$ L to the Topotecan wells. 250  $\mu$ L of media + 0.5% DMSO is added to all remaining wells, into which the test compound(s) are serially diluted. By row, compound-containing media is replica plated (in duplicate) from the assay block to the corresponding cell plates. The cell plates are incubated for 72hours at 37°C, 100% humidity, and 5% CO<sub>2</sub>.

**[0170]      Procedure: Day 4 – MTS Addition and OD Reading:**      The plates are removed from the incubator and 40  $\mu$ L MTS / PMS is added to each well. Plates are then incubated for 120 minutes at 37°C, 100% humidity, 5%CO<sub>2</sub>, followed by reading the ODs at 490nm after a 5 second shaking cycle in a ninety-six well spectrophotometer.


**[0171]      Data Analysis** The normalized % of control (absorbance- background) is calculated and an XLfit is used to generate a dose-response curve from which the concentration of compound required to inhibit viability by 50% is determined.

[0172] The compounds of the present invention show activity when tested in one or more of the methods described in Examples 7, 8 and 9.

[0173] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto. All patents and publications cited above are hereby incorporated by reference.

## WE CLAIM:

1. A compound selected from the group represented by Formula I:



Formula I

where:

U-V is  $-N(R^6)-CR^eR^f-$ ,  $-CR^eR^f-N(R^6)-$ ,  $-N(R^6)-CR^eR^f-CR^gR^h-$ ,  $-CR^eR^f-N(R^6)-CR^gR^h-$  or  $-CR^eR^f-CR^gR^h-N(R^6)-$ ;

$R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are independently hydrogen, alkyl, aryl, aralkyl, heteroaryl, substituted alkyl, substituted aryl, substituted aralkyl or substituted heteroaryl;

$R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are independently hydrogen, alkyl, alkoxy, halogen, cyano or substituted alkyl;

$R^5$  is alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, substituted alkyl, substituted aryl, substituted aralkyl, substituted heteroaryl or substituted heteroaralkyl; and

$R^6$  is hydrogen, acyl, alkyl, aryl, aralkyl, heteroaryl, substituted acyl, substituted alkyl, substituted aryl, substituted aralkyl or substituted heteroaryl;

or a pharmaceutically acceptable salt or solvate thereof.

2. The compound of Claim 1 comprising one or more of the following:

$R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, chloro, fluoro, methyl, methoxy, cyano or substituted lower alkyl;

$R^5$  is aralkyl or substituted aralkyl;

$R^a$  to  $R^h$  are independently hydrogen, lower alkyl or substituted lower alkyl;

U-V is  $-N(R^6)-CR^eR^f-CR^gR^h-$ ,  $-CR^eR^f-N(R^6)-CR^gR^h-$  or  $-CR^eR^f-CR^gR^h-N(R^6)-$ ;

$R^6$  is optionally substituted aralkyl or optionally substituted acyl; and


is an (R)-enantiomer.



3. The compound of Claim 2 comprising one or more of the following:  
R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, chloro, fluoro, methyl, methoxy or cyano;  
R<sup>5</sup> is benzyl or substituted benzyl;  
no more than one of R<sup>a</sup> to R<sup>h</sup> is other than hydrogen;  
U-V is -N(R<sup>6</sup>)-CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>- or -CR<sup>e</sup>R<sup>f</sup>-N(R<sup>6</sup>)-CR<sup>g</sup>R<sup>h</sup>-; and  
R<sup>6</sup> is optionally substituted acyl.
4. The compound of Claim 3 comprising one or more of the following:  
R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen, or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl or cyano;  
R<sup>5</sup> is benzyl;  
R<sup>a</sup> to R<sup>h</sup> are hydrogen;  
U-V is -N(R<sup>6</sup>)-CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-; and  
R<sup>6</sup> is p-methyl-benzoyl.
5. The compound of Claim 4 where: R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> are hydrogen and R<sup>3</sup> is hydrogen or chloro.
6. The compound of Claim 5 where:  
R<sup>5</sup> is benzyl;  
U-V is -N(R<sup>6</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-; and  
R<sup>6</sup> is p-methyl-benzoyl.
7. The compound of Claim 1, selected from:  
3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-piperidin-3-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzyl)-piperidin-4-yl]-3*H*-quinazolin-4-one; and  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-4-yl]-3*H*-quinazolin-4-one.
8. The compound of Claim 7 that is an (R)-enantiomer.

9. The compound of Claim 1, selected from:  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-pyrrolidin-2-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one;  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3*H*-quinazolin-4-one; and  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-4-yl]-3*H*-quinazolin-4-one.
10. The compound of Claim 9 that is an (R)-enantiomer.
11. The compound of Claim 1, selected from:  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-2-yl]-3*H*-quinazolin-4-one; and  
3-benzyl-7-chloro-2-[1-(4-methyl-benzoyl)-piperidin-3-yl]-3*H*-quinazolin-4-one,  
especially the (R)- enantiomers thereof.
12. The compound of Claim 11 that is an (R)-enantiomer.
13. A pharmaceutical formulation comprising a pharmaceutically acceptable excipient and an effective amount of a compound of any of Claims 1-12.
14. A method of treatment comprising administering an effective amount of a compound of any of Claims 1-12 to a patient suffering from a cellular proliferative disease.
15. The method of Claim 14 wherein the cellular proliferative disease is cancer, hyperplasia, restenosis, cardiac hypertrophy, an immune disorder or inflammation.
16. A method of treatment for a cellular proliferative disease comprising administering to a patient suffering therefrom a compound of Claim 1 in an amount sufficient to modulate KSP kinesin activity in cells affected with the disease.
17. A kit comprising a compound of any of Claims 1-12 and a package insert or other labeling including directions for treating a cellular proliferative disease by administering an effective amount of said compound.

18. A compound of the group represented by Formula II:



Formula II

where:

the dashed line indicates that the corresponding bond may be a single bond or a double bond;

T is a covalent bond or optionally substituted lower alkylene;

U-V is chosen from  $-N(R^6)-CR^aR^f-$ ,  $-CR^aR^f-N(R^6)-$ ,  $-N(R^6)-CR^aR^f-CR^gR^h-$ ,  $-CR^aR^f-N(R^6)-CR^gR^h-$ , and  $-CR^aR^f-CR^gR^h-N(R^6)-$ ;

W, X and Y are independently  $-N=$ , N,  $-C=$ , CH,  $CR^i$ , O or S;

Z is  $-N=$ , N,  $-C=$ , CH,  $CR^i$  or is absent, provided that:

no more than two of W, X, Y and Z are  $-N=$ , and

W, X or Y can be O or S only when Z is absent;

$R^i$  is alkyl, alkoxy, halogen, cyano or substituted alkyl;

$R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ ,  $R^e$ ,  $R^f$ ,  $R^g$  and  $R^h$  are independently chosen from hydrogen, alkyl, aryl, aralkyl, heteroaryl, substituted alkyl, substituted aryl, substituted aralkyl and substituted heteroaryl;

$R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are independently chosen from hydrogen, alkyl, alkoxy, halogen, cyano and substituted alkyl;

$R^5$  is alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, substituted alkyl, substituted aryl, substituted aralkyl, substituted heteroaryl or substituted heteroaralkyl; and

$R^6$  is chosen from hydrogen, acyl, alkyl, aryl, aralkyl, heteroaryl, substituted acyl, substituted alkyl, substituted aryl, substituted aralkyl and substituted heteroaryl;

provided that  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  is absent where W, X, Y or Z, respectively, is  $-N=$ , O, S or absent;

or a pharmaceutically acceptable salt or solvate thereof.

19. The compound of Claim 18 comprising one or more of the following:

T is a covalent bond,  $C_1$  to  $C_4$  alkylene or  $C_1$  to  $C_4$  alkylene substituted with halo or oxo;

W, X, Y and Z are independently -C= or -N=;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, chloro, fluoro, methyl, methoxy, cyano or substituted lower alkyl;

R<sup>5</sup> is aralkyl or substituted aralkyl;

R<sup>a</sup> to R<sup>h</sup> are independently hydrogen, lower alkyl or substituted lower alkyl;

U-V is -N(R<sup>6</sup>)-CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-, -CR<sup>e</sup>R<sup>f</sup>-N(R<sup>6</sup>)-CR<sup>g</sup>R<sup>h</sup>- or -CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-N(R<sup>6</sup>)-;

R<sup>6</sup> is optionally substituted aralkyl or optionally substituted acyl; and  
is an (R)-enantiomer.

20. The compound of Claim 19 comprising one or more of the following:

T is a covalent bond or C<sub>1</sub> to C<sub>4</sub> alkylene;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, chloro, fluoro, methyl, methoxy or cyano;

R<sup>5</sup> is benzyl or substituted benzyl;

no more than one of R<sup>a</sup> to R<sup>h</sup> is other than hydrogen;

U-V is -N(R<sup>6</sup>)-CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>- or -CR<sup>e</sup>R<sup>f</sup>-N(R<sup>6</sup>)-CR<sup>g</sup>R<sup>h</sup>-; and

R<sup>6</sup> is optionally substituted acyl.

21. The compound of Claim 20 comprising one or more of the following:

T is a covalent bond;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen, or three of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are hydrogen and the fourth is halo, methoxy, methyl or cyano;

R<sup>5</sup> is benzyl;

R<sup>a</sup> to R<sup>h</sup> are hydrogen;

U-V is -N(R<sup>6</sup>)-CR<sup>e</sup>R<sup>f</sup>-CR<sup>g</sup>R<sup>h</sup>-; and

R<sup>6</sup> is p-methyl-benzoyl.

22. The compound of Claim 21 where:

T is a covalent bond;

R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> are hydrogen and R<sup>3</sup> is hydrogen or chloro;

R<sup>5</sup> is benzyl;

U-V is -N(R<sup>6</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-; and

R<sup>6</sup> is p-methyl-benzoyl.

23. A pharmaceutical formulation comprising a pharmaceutically acceptable excipient and an effective amount of a compound of any of Claims 18-22.
24. A method of treatment comprising administering an effective amount of a compound of any of Claims 18-22 to a patient suffering from a cellular proliferative disease.
25. The method of Claim 24 wherein the cellular proliferative disease is cancer, hyperplasia, restenosis, cardiac hypertrophy, an immune disorder or inflammation.
26. A method of treatment for a cellular proliferative disease comprising administering to a patient suffering therefrom a compound of Claim 18 in an amount sufficient to modulate KSP kinesin activity in cells affected with the disease.
27. A kit comprising a compound of any of Claims 18-22 and a package insert or other labeling including directions for treating a cellular proliferative disease by administering an effective amount of said compound.

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number  
**WO 2004/034972 A3**

(51) International Patent Classification<sup>7</sup>: A01N 43/54,  
A61K 31/517

(21) International Application Number:  
PCT/US2003/030788

(22) International Filing Date:  
30 September 2003 (30.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/414,756 30 September 2002 (30.09.2002) US

(71) Applicant (for all designated States except US): CY-  
TOKINETICS, INC. [US/US]; 280 East Grand Avenue,  
South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BERGNES, Gus-  
tave [US/US]; Cytokinetics, Inc., 280 East Grand Avenue,  
South San Francisco, CA 94080 (US). MORGANS,  
David, J., Jr. [US/US]; 781 Vista Grande Avenue, Los  
Altos, CA 94024 (US).

(74) Agent: GARRET, Arthur, S.; FINNEGAN, HENDER-  
SON, FARABOW, GARRETT & DUNNER, LLP, 1300 I  
Street, N.W., Washington, DC 20005 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
25 November 2004

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 2004/034972 A3

(54) Title: COMPOUNDS, COMPOSITIONS, AND METHODS

(57) Abstract: Compounds, compositions and methods useful for treating cellular proliferative diseases and disorders, for example, by modulating the activity of KSP, are disclosed.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/30788

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 43/54; A61K 31/517

US CL : 514/ 266.22, 266.3

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/ 266.22, 266.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
stic

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,E	Database CAPLUS on STN (Columbus, OH, USA), No. 140:139471, 'Preparation of quinazolinonr-like derivatives to treat cellular proliferative diseases', abstract, WO 2004009036, BERGNES et al. 29 January 2004.	1-27

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 September 2004 (27.09.2004)

Date of mailing of the international search report

12 OCT 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Alton Pryor

Telephone No. 703 308-1235